



Working memory studies among individuals with intellectual disability: An integrative research review



Hefziba Lifshitz ^{a,*}, Esther Kilberg ^b, Eli Vakil ^c

^a School of Education, Bar-Ilan University, Ramat-Gan, Israel

^b Head of Children Development Center, Ramle, Israel

^c Department of Psychology and Leslie and Susan Gonda (Goldschmied) Multidisciplinary Brain Research Centre, Bar-Ilan University, Ramat-Gan, Israel

ARTICLE INFO

Article history:

Received 22 February 2016

Received in revised form 1 August 2016

Accepted 2 August 2016

Available online 7 September 2016

Number of reviews completed is 2

Keywords:

Working memory studies

Research review

Population with NSID

Baddeley and hitch model

Vertical-Horizontal Model

ABSTRACT

Background: Integrative research review infers generalizations about a substantive subject, summarizes the accumulated knowledge that research has left unresolved and generates a new framework on these issues. Due to methodological issues emerging from working memory (WM) studies in the population with non-specific intellectual disability (NSID) ($N = 64$) between 1990–2014, it is difficult to conclude on WM performance in this population.

Aim: This integrative research review aimed to resolve literature conflicts on WM performance among individuals with NSID and to identify the conditions/moderators that govern their WM performance compared to controls with Typical development.

Method/procedure: We used the six stages of integrative research review: problem formulation, data collection, evaluation, data analysis, results, interpretation and discussion.

Outcomes and results: The findings indicate two types of moderators that determine WM performance in the population with NSID: Participants' moderators (criteria for matching the ID and TD groups, CA and MA), and **task** moderators [the three WM components of Baddeley and Hitch's (1974) model and task load]. Only an interaction between the two moderators determines WM performance in this population. The findings indicate a hierarchy (from more to less preserved) in WM performance of individuals with NSID: The visuospatial tasks, then some of the executive functions tasks, and the phonological loop tasks being less preserved. Furthermore, at a low level of control, the performance of participants with NSID was preserved beyond the modality and vice versa.

Conclusions and implications: Modality and MA/intelligence determine WM performance of individuals with ID. Educators should prepare intervention programs take the impact of the two moderators into account.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

“Working memory refers to a broad framework of interacting processes that involve the temporary storage and manipulation of information in the service of performing complex cognitive activities” (Baddeley, Allen, & Hitch, 2011, p. 1393).

* Corresponding author.
E-mail addresses: hefziba@013net.net, Hefziba.lifshitz@biu.ac.il (H. Lifshitz).

It evolved from the earlier concept of short-term memory (STM), from which it differs by emphasizing the functional importance of the system and by replacing a unitary store with a multicomponent system (Baddeley & Hitch, 1974).

Scientists are in disagreement on whether WM per se explains fluid intelligence (Garlick & Sejnowski, 2006) and whether the working memory (WM) capacity is associated with fluid intelligence (g) (Conway et al., 2005; Garlick & Sejnowski, 2006; Kane & Engle 2002). Baddeley (2000) (and in Baddeley et al., 2011) claimed that "Attention and temporary storage, are themselves unchanged by learning, other than indirectly via the crystallized systems". Kane and Engle (2002) stated that the "Executive attention, is the psychological core of the statistical construct of general fluid intelligence, or psychometric Gf (page 638)".

It has been documented that WM is associated with cognitive skills such as problem solving, reasoning and academic achievements (Bayliss, Jarrold, Baddeley, & Leigh, 2005; Hitch & McAuley, 1991) as well as with everyday life skills such as reading, writing, arithmetic, and language (Gathercole, Alloway, Willis, & Adams, 2004; Numminen, Service, & Ruoppila, 2002) which enable an independent life. Thus, the importance of WM in the population with intellectual disability) is thus self-evident (Van der Molen, Luit, Van de Molen, & Jengmans, 2011).

The most widely recognized model of WM is that of Baddeley and Hitch (1974) and Baddeley (2003, 2008). This model assumes a limited capacity controller, the central executive, supported by two temporary slave systems, the phonological loop and the visuospatial sketchpad. The phonological loop involves the temporary storage of phonological auditory information and is comprised of two sub-systems: The phonological store and the sub-vocal rehearsal process (Schuchardt, Maehler, & Hasselhorn, 2011). The visuospatial sketchpad is assumed to be capable of holding and manipulating visual and spatial information. The use of spatial imagery in immediate recall is disrupted by tasks such as tracking a moving object (Baddeley, Thomson, & Buchanan, 1975), while memory for pattern and shape is disrupted by the passive processing of line drawings, or even colour patches (Logie & Pearson, 1997; Logie, 1986). The central executive is involved in conditions of high level processing, including a set of high level attentional cognitive abilities such as planning, attention, inhibition and shifting (Baddeley et al., 2011; Camos & Barrouillet, 2014). A neutral modality storage, the episodic buffer, has recently been proposed (Baddeley et al., 2011), which is assumed to be a temporary multidimensional store that forms an interface between the subsystems of WM, long-term memory (LTM) and the central executive. Its major function is to bind different sources of information into integrated chunks (Baddeley et al., 2011). The phonological loop, the visuospatial sketchpad and the central executive were examined broadly in the population with ID, whereas the episodic buffer was examined in only one study (Henry, 2010), and will therefore not be included in the present review.

Cornoldi, Rigoni, Venneri, and Vecchi (2000) argued that Baddeley's model cannot by itself explain WM performance patterns of individuals with ID. Cornoldi and Vecchi (2003) suggested a double '*Horizontal and vertical continuum*' model. The *horizontal continuum* relates to the above-mentioned components of Baddeley et al.'s (2011) model. The *vertical continuum* reflects the required degree of control which is defined by the amount of active processing necessary for manipulating the information maintained in a temporary memory system ranging from passive to active tasks (Cornoldi & Vecchi, 2003), i.e. the cognitive load inherent in the task. Some tasks that measure the articulatory loop or visuospatial sketchpad simply require remembering the material as it was presented, and involve a low level of control, while other tasks require a high level of control. The concept of cognitive load raised by Cornoldi and Vecchi (2003) parallels the concept of attention and cognitive load suggested by Camos and Barrouillet (2014). They define cognitive load as the duration of attentional capture divided by the total time of performing the task.

Scientists are interested in the question of whether WM is domain-specific or domain-general (Baddeley & Logie, 1999). In their time-based resource-sharing model, Camos and Barrouillet (2014) argue that domain-general resources are responsible for processing and storage and that attention is involved in both. They claim two systems that are involved in the maintenance of verbal information: the phonological loop and the executive loop. The phonological loop is less attention-demanding (primary memory according to Unsworth & Engel, 2007) and involves the recalling of simple verbal tasks (i.e., digit span forward). The executive loop involves manipulation of the task in addition to just recalling, and demands a higher level of attention according to the tasks (inhibition, selecting, etc.).

Use of the phonological loop makes recall sensitive to the phonological characteristics of the material to be maintained, whereas the phonological nature of the memory items does not affect recall performance under the use of the executive loop. This does not imply that the executive loop is a "better" system of maintenance that should always be favored for verbal information. Because attentional refreshing is more attention-demanding than sub-vocal rehearsal, the former is very sensitive to the availability of attention and the presence of concurrent attention-demanding tasks. Attention is involved in both the retention of information and in processing activities. The capacity of the focus of attention appears clearly when verbal rehearsal is prevented, suggesting that maintenance of verbal information through attentional focusing and verbal rehearsal must be distinguished. Brain imaging studies reported distinct neural structures supporting the separation of these two loops.

As far as we know, only one study tapped the effect of cognitive load on WM performance among individuals with Down syndrome versus peers with typical development (TD) with the same mental age (MA) (Lanfranchi, Cornoldi, & Vianello, 2004). It was found that in tasks requiring a low level of control, children with Down syndrome showed impairment of verbal but not visuospatial WM tasks. As the requirement for control increased, they showed greater impairment on both tasks. The above study implies the need to examine the effect of cognitive load in WM tasks among individuals with non-specific

ID (NSID). In this integrative review we examined the effect of the level of control on WM performance among individuals with NSID beyond modality.

1.1. Cognitive profile of individuals with NSID

Individuals with ID are a heterogeneous group in their IQ level, etiology and associated disorders. Poor language competence (Fink & Cegelka, 1982), lack of verbal rehearsal and reduced ability to retrieve stored information (Hulme & Mackenzie, 1992) are common. They exhibit lack of cognitive strategies in dealing with problems, as well as in the use of meta-cognitive strategies (Borkowski, Carr, & Pressley, 1987), difficulties in shifting flexibly from one strategy to another (Campione & Brown, 1984), attention deficit (Reed, 1996), and lack of automatic identification of presented stimuli (Ellis, 1978). They also exhibit inefficient STM (Ellis, 1978). Some of the studies found equal WM performance in the population with ID compared to participants with TD (Danielsson, Henry, Rönnberg, & Nilsson, 2010; Henry, 2001; Van der Molen, Van Luit, Jongmans, & Van der Molen, 2007), whereas others found lower WM performance (Alloway, 2010; Conners, Carr, & Willis, 1998; Hartman, Houwen, Scherder, & Visscher, 2010; Hasselhorn & Mähler, 2007; Henry & MacLean, 2002; Van der Molen, Van Luit, Jongmans, & Van der Molen, 2009; Van der Molen, Van Luit, Van der Molen, & Jongmans, 2010; Willner, Bailey, Parry, & Dymond, 2010). The question is whether there is a general “law” which dictates the equal or lower performance of the population with ID compared to the population with TD.

1.2. Integrative research review

“Integrative research review infers generalizations about a substantive subject from a set of studies bearing on the same issues” (Cooper, 1982, p. 438; Randolph, 2009; Torraco, 2005). “Its goal is to summarize the accumulated knowledge regarding the relation (s) of interest, highlight important issues, critiques, synthesize representative literature that research has left unresolved and generate a new framework on these issues” (Torraco, 2005; p. 3). Several methodological issues emerge from the WM studies we reviewed, which make it difficult to draw definitive conclusions. Some focus on the phonological loop, some on the visuospatial sketchpad and others on executive function tasks. Furthermore, different intelligence tests were used for matching between groups. There is also a lack of distinction between ID levels, and different WM tests were used for measuring the same skills. The goal of this review was to resolve literature conflicts on WM performance among individuals with ID and to identify the conditions/moderators that govern their WM performance compared to controls with TD.

One might argue that conducting a meta-analysis on WM studies in the population with ID is the best method for answering this question. Lifshitz, Stein, Weiss and Vakil (2011) published a meta-analysis of 40 explicit memory studies of three etiologies with ID. The explicit memory of individuals with ID was found to be poorer than in controls with TD with the same chronological age (CA) or mental age (MA). Relatively preserved explicit memory performance was found among individuals with Williams syndrome compared to Down syndrome. Cohen (1988, 1992) argued that a ‘summary effect’ could mask findings that are insufficient to alter the generalizations, but might hold important qualitative data that are potentially significant for rehabilitational and educational purposes. The meta-analysis revealed poorer performance in recall and recognition in verbal and visual tasks. However, examining each study separately revealed that recognition and free recall in the visual modality were preserved in some of the studies (Dulaney & Ellis, 1991; Perrig & Perrig, 1995). Using a different methodology – an integrative research review (Lifshitz-Vahav & Vakil, 2014), we elucidated the underlying moderators of preserved/impaired explicit memory studies. We suggested a taxonomy of moderators related to the participants and the tasks, where only an interrelation between the two dictates memory performance in the population with ID.

We shall review WM studies in the population with NSID from 1990. Until that time, memory studies in the population with ID focused only on STM (Atkinson & Shiffrin, 1968, 1971; Baddeley, 2000). The concept of WM was introduced to the field of ID only in the 1990s. The theoretical foundations of this integrative review were the vertical and horizontal WM models (Cornoldi & Vecchi, 2003). The **horizontal** model relates to the components of Baddeley’s model: The phonological loop, the visuospatial sketchpad, and the executive function, whereas the **vertical** model relates to the level of control required in the task from passive to active manipulation in each component (Baddeley et al., 2011).

The operative goals were: (a) To find the WM phenotype in each component of Baddeley’s model among participants with NSID. Put differently, we aimed to find the moderators that influence WM performance in Baddeley’s model. Lifshitz and Vakil (2014) found the “picture superiority effect” in explicit memory studies among individuals with ID. That is, none of the verbal explicit tasks yielded equal performance of individuals with ID compared to those with TD. Based on the above we hypothesized that participants with NSID would exhibit relatively more preserved visual WM tasks compared to phonological loop tasks. (b) To examine the effect of cognitive load in WM tasks beyond modality. We hypothesized that WM performance in the population with NSID would vary according to the cognitive load of the task. Performance would decrease with increasing cognitive load.

2. Method

We conducted a computerized search of articles on ID published between 1990 and 2014 in journals that focus on memory and cognition and in journals that focus on ID, using the PsycNET database, Dissertation Abstracts International, the




Fig. 1. Taxonomy of moderators that influence working memory in a population with ID.

Social Science Citation Index, ERIC, Webspires, Proquest, Ebsco, and Google Scholar. Keywords were “short term memory,” “working memory,” “phonological loop,” “visuospatial sketchpad,” “executive function,” “mental retardation” or “intellectual disability”; and “non-specific ID or etiology.” Inclusion criteria were: studies of individuals with ID without other disabilities such as autism or mental illness; IQ ranging from mild to moderate ID; samples larger than a case study; a control group of individuals with TD; studies that included statistical analyses or presented means and standard deviations.

The search yielded 64 studies (157 experiments) of WM in the population with NSID, which were conducted between 1990 and 2014. Only 15% ($n=10$) were conducted in 1990–1999, while 85% ($n=54$) were conducted in 2001–2011. Thus, WM research in this population is still relatively young.

Some of the studies conducted several experiments in the same study using different modalities and tests. Each of these was considered an experiment. Table 1 presents the distribution of WM experiments in the population with NSID according to Baddeley and Hitch (1974) and Baddeley et al.'s (2011) model.

The following conclusions emerge from the table: 15% ($n=10$) of the studies focused on children aged 5–12; 70% ($n=44$) on adolescents aged 12–21; and only 15% ($n=10$) on adults aged 21–50. Thus, adolescents are currently at the focus of WM studies in the population with NSID. 43% ($n=68$) of the experiments focused on the phonological loop; 21% ($n=32$) on the visuospatial sketchpad; and 37% ($n=57$) on the central executive.

2.1. Procedure

An integrative research review includes six stages (Cooper, 1982; Randolph, 2009; Torraco, 2005): (1) problem formulation; (2) data collection; (3) data evaluation; (4) data analysis; (5) presentation of results; and (6) interpretation and discussion. We used a double factor model that we used previously for examining explicit LTM studies in the population with ID (Lifshitz & Vakil, 2014), albeit with adaptation to WM studies: Moderators related to **background characteristics of participants** (criteria for matching the ID and TD groups, CA and MA), and moderators related to the **task** (the three components of Baddeley et al.'s (2011) model: the phonological loop, the visuospatial sketchpad and executive function). The tentative moderators used in this integrative review are presented in Fig. 1. We triangulated (Randolph, 2009) between the moderators in order to find their involvement when memory was preserved or impaired. In cases of contradictory results for the same moderators, we re-read the articles to find other moderators that might influence performance. In addition, we counted the number of preserved/impaired experiments in each component. Based on the above, we constructed a table which relates to the hierarchy of the WM tasks in the components in Baddeley's model (Baddeley & Hitch, 1974) (from more to less preserved). It should be noted that the content of the tasks is different in each component, but we compared the number of preserved/impaired performances in each component beyond type of task. Furthermore, this is an integrative research review which is a kind of qualitative study (Randolph, 2009; Torraco, 2005) with different goals than a meta-analysis. As stated earlier (p. 7), a ‘summary effect’ could ignore findings that might hold important qualitative data that are potentially significant for rehabilitation and educational purposes (Cohen, 1988; 1992). Therefore, our WM hierarchy tasks table does not take the sample size and effect size into account, but is based on percentage which was found to be a reliable measure in our previous integrative research review (Lifshitz-Vahav & Vakil, 2014). We then attempted to integrate the results. In the discussion we interpreted the results and placed them in a theoretical framework.

2.2. Findings

Table 1 presents the results of the three components of WM among individuals with NSID. **Table 2** relates to the hierarchy of the WM tasks in the three components. The findings will be presented according to the three components, and moderators related to both participants and tasks will be discussed.

Table 1

Phonological loop, visuospatial span and central executive experiments among participants with NSID, compared to participants with TD matched by CA, MA and/or IQ: tests and findings (Alloway, Gathercole, & Pickering, 2007; Baddeley, 1997; Belacchi, Carretti, & Cornoldi, 2010; Bellugi, Lichtenberger, Jones, Lai, & St George, 2000; Danielsson, Henry, Messer, & Rönnberg, 2012; Gathercole, 1999; Gathercole, Adams, & Hitch, 1994; Gathercole & Alloway, 2006; Gunn & Crombie, 1996; Hulme, Maughan, & Brown, 1991; Lanfranchi, Carretti et al., 2009; Lanfranchi, Cornoldi et al., 2009; Laws, 2002; Levén, Lyxell, Andersson, Danielsson, & Rönnberg, 2008; Levén, Lyxell, Andersson, Danielsson, & Rönnberg, 2011; Maehler & Schuchardt, 2009; McCarthy, 1972; Merighi, Edison, & Zigler, 1990; Miller, 1992; Miyake, Friedman, Rettinger, Shah, & Hegarty, 2001; Nash & Heath, 2011; Numminen, Service, Ahonen, & Ruoppila, 2001; O'Hearn, Courtney, Street, & Landau, 2009; Pazzaglia & Cornoldi, 1999; Piaget, 1970; Pickering & Gathercole, 2001; Poloczek, Büttner, & Hasselhorn, 2014; Porter, Coltheart, & Langdon, 2007; Rondal, 1996; Schuchardt, Gebhardt, & Mäehler, 2010; Vicari, Bellucci, & Carlesimo, 2003; Vicari, Bellucci, & Carlesimo, 2005; Vicari, Carlesimo, & Caltagirone, 1995; Wang & Bellugi, 1994; Wechsler, 1967).

| Article name | Group | CA | MA | IQ | Working measures/The test and the modality | Results |
|--|--------------------------|------------------|-------|---|---|---|
| Digit | | | | | | |
| Leken et al. (2011) | Mild ID (<i>n</i> = 10) | 275 | | 85-11 | Digit forward | Impaired |
| | TD (<i>n</i> = 5) | 27 | | 50-70 | | |
| Numminen et al. (2002) | NSID (<i>n</i> = 24) | 49.88 | | Raven | Digit Span | |
| | TDMA (<i>n</i> = 24) | 5.650 (3-6) | | Equal fluid intelligence | Forward Task | preserved |
| Numminen et al. (2001) | NSID (<i>n</i> = 24) | 38-58 (49.88) | 5.63 | Raven | Digit span forward | Preserved |
| | TDMA (<i>n</i> = 24) | 3-6 (5.50) | 5.67 | | | |
| Conners, Carr, and Willis (1998) | NSID (<i>n</i> = 30) | 11.72 | 7.45 | 66.58 (WISC-R verbal) | Forward digit span | Impaired CA Preserved MA CA>ID=MA |
| | TDCA (<i>n</i> = 30) | | | 107.42 | | |
| | TDVMA (<i>n</i> = 26) | 12.36 | | 106.77 | | |
| | | 7.45 | 7.45 | | | |
| Vicari et al. (1995) | DS (<i>n</i> = 15) | 16.6 | 5.3 | 36 Leiter | Forward digit span | Preserved |
| | NSID (<i>n</i> = 14) | 16.7 | 51 | 39 | | |
| | TDMA (<i>n</i> = 24) | 5.7 | 5.5 | 101 | | |
| Henry and Winfield (2010) | MID (<i>n</i> = 35) | 12-13 (12.6) | 7.7 | 57 | digit span | Impaired |
| | TDMA (<i>n</i> = 35) | 6-8 (7.5) | 7.8 | 101 | | |
| Schuchardt, Gebhardt, and Maehler (2010) | BID (<i>n</i> = 15) | 15 | | British Ability S. | | |
| | MID (<i>n</i> = 15) | 15 | | 82.63 (70-84) 61.82 (50-69) (K_ABC) | Memory span for digit | Impaired |
| | TDCA (<i>n</i> = 25) | 15 | | 109.80 (90-115) | | |
| | BID (<i>n</i> = 19) | 10 | | 81.21 (CMMS) | | |
| | TDMA (<i>n</i> = 22) | 7 | | 104.91 | | |
| Van der Molen et al. (2010) | Mild ID (<i>n</i> = 39) | 13-17 | 118.1 | 65 (55-75) (WISC_R) | Digit Recall (Baddeley, 2008) | Preserved MA Impaired Ca |
| | TDCA (<i>n</i> = 39) | 13-16 | 182.6 | 99.8 | | |
| | TDMA (<i>n</i> = 26) | 9-12 | 120.7 | 97.1 | | |
| Maehler and Schuchardt (2009) | NSID (<i>n</i> = 27) | 12.45 | | 75.44 | Digit span | Impaired |
| | LD (<i>n</i> = 27) | 12.35 | | 100.08 | | ID=LD<TD |
| | TDCA (<i>n</i> = 27) | 10.58 | | 101.04 | | |
| Van der Molen et al. (2009) | Mild ID (<i>n</i> = 49) | 13-17 | | WISC-III | Digit Recall | Impaired CA Preserved MA |
| | TDCA (<i>n</i> = 39) | 13-16 | | 55-85 | | |
| | TDMA (<i>n</i> = 29) | 8-12 | | | | |
| Van der Molen et al. (2007) | Mild ID (<i>n</i> = 50) | 15.3 | | Raven | Digit span | Impaired CA and MA |
| | TDCA (<i>n</i> = 25) | 15.3 | | 55-85 | | |
| | TDMA (<i>n</i> = 25) | 10.10 | | | | |
| Bayliss et al. (2005) | IDvLD (<i>n</i> = 50) | 13.6 (10.1-16.3) | 7.5 | 55.32 (27.72-69.89) | Digit span | Impaired |
| | TDVMA (<i>n</i> = 50) | 7.3 (5.5-9.1) | 7.5 | 102.36 (80.81-129.17) BPVS-II | | |
| Henry and MacLean (2003) | NSID (<i>n</i> = 53) | 11-12(11.11) | | 57.2 (40-79) | Digit span | Preserved |
| | TDCA (<i>n</i> = 45) | 11-12 (12.1) | | 103.8 (84-123) | | |
| | TDMA (<i>n</i> = 41) | 7-8 (7.11) | | 101.0 (82-136) | | |
| Henry and MacLean (2002) | NSID (<i>n</i> = 53) | 11-12 | | 57.2 (BAS) | Digit span | Preserved MA Impaired CA |
| | TDCA (<i>n</i> = 45) | 11-12 | | 103.8 | | |
| | TDMA (<i>n</i> = 41) | 7-8 | | 101.0 | | |
| Henry (2001) | BID (<i>n</i> = 10) | 11-12 | 9.4 | 75.3 (70-79) | Digit span | Impaired |
| | Mild ID (<i>n</i> = 25) | 11-12 | 7.9 | 60.5 (55-68) | | |
| | MI D (<i>n</i> = 21) | 11-12 | 6.4 | 45.8 (40-52) | | |
| | TD CA (<i>n</i> = 22) | 11-12 | 13.0 | 104.5 (84-123) | | |
| Hulme et al. (1991) | NSID (<i>n</i> = 55) | 19.1 (9.9-38.2) | 6.3 | 20-50 | ITPA auditory sequential memory test – digit span | Impaired |
| | TDMA (<i>n</i> = 55) | 6.0 (4.2-8.4) | 6.0 | | Longitudinal study after 2 and 5 years | |

Table 1 (Continued)

| Words | | | | | | | |
|---|--------------------------|---------------|------|--------------------------|--|---|-------------------------------------|
| Schuchardt et al. (2011) | BID (<i>n</i> = 19) | 10 | 7 | 80.0 (70-84) CMM) | Word span | Mild ID impaired in phonological storage | |
| | Mild ID (<i>n</i> = 22) | 15 | 7 | 61.82 (50-69) (K_ABC) | One-syllable dissimilar | BID = TDMA | |
| | TDMA (<i>n</i> = 22) | 7 | 7 | 106.73 (93-115) (CMM) | | | |
| Carretti et al. (2010) | NSID (<i>n</i> = 28) | 38.4 | 6.2 | Raven:17.36 (IQ = 40-75) | <u>Verbal WM tests</u> | | |
| | TDMA (<i>n</i> = 28) | 6.5 | 6.6 | 17.50 | (Belacchi, Carretti, & Cornoldi, 2010) | Preserved | |
| Henry and Winfield (2010) | MID (<i>n</i> = 35) | 12-13 (12.6) | 7.7 | 57 (BAS) | Forward word span | Impaired | |
| | TDMA (<i>n</i> = 35) | 6-8 (7.5) | 7.8 | 101 | Word span picture span | Impaired | |
| Mahler and Schuchardt (2009) | NSID (<i>n</i> = 27) | 12.45 | | 75.44 | Word span (one-three syllable words) | Impaired | |
| | LD (<i>n</i> = 27) | 12.35 | | 100.08 | ID=LD<TD | | |
| | TDCA (<i>n</i> = 27) | 10.58 | | 101.04 | | | |
| Hasselhorn (2007) | NSID (<i>n</i> = 22) | 10.7 | | 78.9 (CMM 1-4) | Word span | Impaired CA | |
| | TDCA (<i>n</i> = 22) | 10.5 | 7.5 | 100.5 | Preserved MA | | |
| | TDMA (<i>n</i> = 22) | 7.4 | | 107.8 | | | |
| Henry and MacLean (2003) | NSID (<i>n</i> = 53) | 11-12 (11.11) | | 57.2 (40-79) (BAS) | Word span | Impaired | |
| | TDCA (<i>n</i> = 45) | 11-12 (12.1) | | 103.8 (84-123) | Word span | | |
| | TDMA (<i>n</i> = 41) | 7-8 (7.11) | | 101.0 (82-136) | | | |
| Henry and MacLean (2002) | NSID (<i>n</i> = 53) | 11-12 | | 57.2 (BAS) | Word span | Impaired | |
| | TDCA (<i>n</i> = 45) | 11-12 | | 103.8 | 101.0 | | |
| | TDMA (<i>n</i> = 41) | 7-8 | | | | | |
| Lanfranchi et al. (2002) | DS (<i>n</i> = 17) | 8-16 (11.11) | 5.6 | Logical Operational Test | <u>Low control-</u> | Preserved | |
| | NSID (<i>n</i> = 13) | 8-16 | 5.6 | | Forward word span | DS=NSID=TD | |
| | TDMA (<i>n</i> = 30) | 5.2 | 5.6 | | | | |
| Henry (2001) | BID (<i>n</i> = 10) | 11-12 | 9.4 | 75.3 (70-79) | Word span | Impaired | |
| | Mild ID (<i>n</i> = 25) | 11-12 | 7.9 | 60.5 (55-68) | 104.5 (84-123) | | |
| | MI D (<i>n</i> = 21) | 11-12 | 6.4 | 45.8 (40-52) | | | |
| Hulme and Mackenzie (1992) | TD CA (<i>n</i> = 22) | 11-12 | 13.0 | | | | |
| | | | | BAS | | | |
| | NSID (<i>n</i> = 24) | 13.4 | 14.6 | 5.2 5.9 6.10 | EPVT | Serial recall | Impaired |
| | TDMA (<i>n</i> = 24) | 5.2 | 6.0 | 6.9 | 5.2 5.10 6.11 | 20-50 | |
| Word Length Effect | | | | | | | |
| Bordeline | Mild ID (<i>n</i> = 34) | 11 | | 72 | Word length effect | Impaired | |
| | TD (<i>n</i> = 34) | 7.6 | | 94 | | | |
| Schuchardt et al. (2011) | BID (<i>n</i> = 19) | 10 | | 80.0 (70-84) (CMM) | One-Three syllables dissimilar | Preserved -All three groups show word length effect | |
| | Mild ID (<i>n</i> = 22) | 15 | | 61.82 (50-69) (K_ABC) | | | |
| | TDMA (<i>n</i> = 22) | 7 | | 106.73 (93-115) (CMM) | | | |
| Hasselhorn (2007) | NSID (<i>n</i> = 22) | 10.7 | | 78.9 (54-88) | Memory span – word length effect | Impaired CA | |
| | TDCA (<i>n</i> = 22) | 10.5 | | 100.5 | (1.3 syllables words) | Preserved MA | |
| | TDMA (<i>n</i> = 22) | 7.4 | | 107.8 | | | |
| Van der Molen et al. (2007) | Mild ID (<i>n</i> = 50) | 15.3 | | Raven | Non-word test | Impaired CA | |
| | TDCA (<i>n</i> = 25) | 15.3 | | 55-85 | (monosyllabic vs. two syllabic words) | Preserved MA | |
| | TDMA (<i>n</i> = 25) | 10.10 | | | | | |
| Rosenquist, Conners, and Roskos-Ewoldsen (2003) | NSID (<i>n</i> = 19) | 12-16 | 8.08 | 55-70 (63.3) WISC-R | Word length task | -No difference in ID between short and long words | |
| | TDMA (<i>n</i> = 23) | 7-8 | 8.21 | | | | |
| | | | | | | | |
| Hulme and Mackenzie (1992) | NSID (<i>n</i> = 24) | 5.2 | 6.0 | 6.9 | EPVT | Serial recall – word length (one, two & three syllable words) | Impaired ID Non word length effects |
| | TDMA (<i>n</i> = 24) | 5.2 | 5.10 | 6.11 | 20-50 | | |
| | | | | | | | |
| Similarity effect | | | | | | | |
| Schuchardt et al. (2011) | BID (<i>n</i> = 19) | 10 | | 80.0 (70-84) (CMM) | One-syllable dissimilar/ dissimilar | Preserved -All three groups show similarity effect | |
| | Mild ID (<i>n</i> = 22) | 15 | | 61.82 (50-69) (K_ABC) | | | |
| | TDMA (<i>n</i> = 22) | 7 | | 106.73 (93-115) (CMM) | | | |

Table 1 (Continued)

| | | | | | |
|-------------------------------|--|---|---|---|--|
| Hasselhorn (2007) | NSID (<i>n</i> = 22) TDCA (<i>n</i> = 22) TDMA (<i>n</i> = 22) | 10.7 10.5 7.4 | 78.9 (54-88) 100.5 107.8 (CMM 1-4) | Memory span – phonological similarity effect (one syllables similar/dissimilar) | Preserved – similarity in the phonological similarity effect |
| Rosenquist et al. (2003) | NSID (<i>n</i> = 19) TDMA (<i>n</i> = 23) | 12-16 7-8 | 8.08 8.21 | 55-70 (63.3) | Phonological similarity test (Hulme et al., 1991) |
| Hulme and Mackenzie (1992) | NSID (<i>n</i> = 24) T DMA (<i>n</i> = 24) | 13.4 14.6 16.5 5.2 6.0 6.9 | 5.2 5.9 6.10 5.2 5.10 6.11 | EPVT | Memory span test for acoustically similar and dissimilar words |
| Non-Words | | | | | |
| Schuchardt et al. (2011) | BID (<i>n</i> = 19) Mild ID (<i>n</i> = 22) TDMA (<i>n</i> = 22) | 10 15 7 | 7 7 7 | 80.0 (70-84) (CMM) 61.82 (50-69) (K_ABC) 106.73 (93-115) (CMM) | Mild ID impaired in phonological storage BID = TDMA |
| Schuchardt et al. (2010) | BID (<i>n</i> = 15) MID (<i>n</i> = 15) TDCA (<i>n</i> = 25) BID (<i>n</i> = 19) TDMA (<i>n</i> = 22) | 15 15 15 10 7 | 7.1 7.1 7.1 7.1 7.1 | 82.63 (70-84) 61.82 (50-69) 109.80 (90-115) 81.21 104.91 | One-syllable non-words Non-word repetition |
| Van der Molen et al. (2010) | Mild ID (<i>n</i> = 39) TDCA (<i>n</i> = 39) TDMA (<i>n</i> = 26) | 13-17 13-16 9-12 | 118.1 182.6 120.7 | 65 (55-75) (WISC-III) 99.8 97.1 | Non word recall |
| Maehler and Schuchardt (2009) | NSID (<i>n</i> = 27) LD (<i>n</i> = 27) TDCA (<i>n</i> = 27) | 12.45 12.35 10.58 | | 75.44 100.08 101.04 | Non-word repetition ID=LD<TD |
| Van der Molen et al. (2009) | Mild ID (<i>n</i> = 49) TDCA (<i>n</i> = 39) TDMA (<i>n</i> = 29) | 13-17 13-16 8-12 | | 55-85 (WISC-III) | Non Word Recall Impaired MA and CA |
| Hasselhorn (2007) | NSID (<i>n</i> = 22) TDCA (<i>n</i> = 22) TDMA (<i>n</i> = 22) | 10.7 10.5 7.4 | 7.5 | 78.9 (CMM 1-4) 100.5 107.8 | Non word repetition (2-4 syllables) |
| Van der Molen et al. (2007) | Mild ID (<i>n</i> = 50) TDCA (<i>n</i> = 25) TDMA (<i>n</i> = 25) | 15.3 15.3 10.10 | | Raven 55-85 | Non word test Impaired CA Impaired MA |
| Numminen et al. (2002) | NSID (<i>n</i> = 24) TDMA (<i>n</i> = 24) | 49.88 5.650 (3-6) | 5.63 5.67 | Raven Equal fluid intelligence Non Word Repetition task Non Word Span Task Finish non word task | Non Word Repetition task Non Word Span Task Impaired |
| Numminen et al. (2001) | NSID (<i>n</i> = 24) TDMA (<i>n</i> = 24) | 38-58 (49.88) 3-6 (5.50) | | Raven | Finish non word task Impaired |
| Visual-Object | | | | | |
| Henry and Winfield (2010) | MID (<i>n</i> = 35) TDMA (<i>n</i> = 35) | 12-13 (12.6) 6-8 (7.5) | 7.7 7.8 | 57 101 | Pattern span Preserved in same mental age |
| Van der Molen et al. (2010) | Mild ID (<i>n</i> = 39) TDCA (<i>n</i> = 39) TDMA (<i>n</i> = 26) | 13-17 13-16 9-12 | 118.1 182.6 120.7 | 65 (55-75) 99.8 97.1 | Visual pattern tests (Della Sala, 1997) Preserved MA Impaired CA |
| Maehler and Schuchardt (2009) | NSID (<i>n</i> = 27) LD (<i>n</i> = 27) TD (<i>n</i> = 27) | 12.45 12.35 10.58 | | 75.44 100.08 101.04 | matrix span simple matrix span complex |
| Van der Molen et al. (2009) | Mild ID (<i>n</i> = 49) TDCA (<i>n</i> = 39) TDMA (<i>n</i> = 29) | 13-17 13-16 8-12 | | | Visual patterns test Impaired CA preserved MA |
| Henry and MacLean (2003) | NSID (<i>n</i> = 53) TDCA (<i>n</i> = 45) TDMA (<i>n</i> = 41) | 11-12 (11.11) 11-12 (12.1) 7-8 (7.11) | | 57.2 (40-79) 103.8 (84-123) 101.0 (82-136) | Pattern span Impaired CA Preserved MA |
| Rosenquist et al. (2003) | NSID (<i>n</i> = 19) TDMA (<i>n</i> = 23) | 12-16 7-8 | 8.08 8.21 | 55-70 (63.3) | Visual complexity task Visual similarity task |
| Henry and MacLean (2002) | NSID (<i>n</i> = 53) TDCA (<i>n</i> = 45) TDMA (<i>n</i> = 41) | 11-12 11-12 7-8 | | 57.2 BAS 103.8 101.0 | Pattern span Preserved MA ID>MA Impaired CA |
| Henry (2001) | BID (<i>n</i> = 10) Mild ID (<i>n</i> = 25) | 11-12 11-12 | 9.4 7.9 | 75.3 (70-79) 60.5 (55-68) | Pattern span Impaired MIQ & mild ID |

Table 1 (Continued)

| | MI D (n = 21) TD CA (n = 22) | 11-12 11-12 | 6.4 13.0 | 45.8 (40-52) 104.5 (84-123) | Preserved BIQ |
|---------------------------------------|--|---|---------------------------|--|---|
| Visual-Spatial | | | | | |
| Henry and Winfield (2010) | MID (n = 35) TDMA (n = 35) | 12-13 (12.6) 6-8 (7.5) | 7.7 7.8 | 57 101 | Spatial span Preserved in same mental age |
| Schuchardt et al. (2010) | BID (n = 15) MID (n = 15) TDCA (n = 25) BID (n = 19) TDMA (n = 22) | 15 15 15 10 7 | | 82.63 (70-84) 61.82 (50-69) 109.80 (90-115) 81.21 104.91 | Memory span for locations Corsi block simple Corsi block complex Preserved in the same mental age |
| Van der Molen et al. (2010) | Mild ID (n = 39) TDCA (n = 39) TDMA (n = 26) | 13-17 13-16 9-12 | 118.1 182.6 120.7 | 65 (55-75) 99.8 97.1 | Block recall=Corsi Impaired |
| Maehler and Schuchardt (2009) | NSID (n = 27) LD (n = 27) TD (n = 27) | 12.45 12.35 10.58 | | 75.44 100.08 101.04 | Memory span for locations Corsi-block simple Corsi block complex Preserved Impaired ID=LD<TD |
| Van der Molen et al. (2009) | Mild ID (n = 49) TDCA (n = 39) TDMA (n = 29) | 13-17 13-16 8-12 | | | Corsi blocks task Impaired MA & CA |
| Bayliss et al. (2005) | IDLD (n = 50) TDVMA (n = 50) | 13.6 (9.10-16.3) 7.3 (5.5-9.1) | 7.5 7.5 | 55.32 (27.72-69.89) BPVS-II 102.36 (80.81-129.17) | Corsi span Impaired |
| Henry and MacLean (2003) | NSID (n = 53) TDCA (n = 45) TDMA (n = 41) | 11-12 (11.11) 11-12 (12.1) 7-8 (7.11) | | 57.2 (40-79) 103.8 (84-123) 101.0 (82-136) | Visuo-spatial span preserved |
| Henry and MacLean (2002) | NSID (n = 53) TDCA (n = 45) TDMA (n = 41) | 11-12 11-12 7-8 | | 57.2 BAS 103.8 101.0 | Spatial span (Corsi) Preserved MA ID>MA |
| Numminen et al. (2002) | NSID (n = 24) TDMA (n = 24) | 49.88 5.650 (3-6) | | Raven | Corsi Blocks Task Impaired |
| Henry (2001) | BID (n = 10) Mild ID (n = 25) MI D (n = 21) TD CA (n = 22) | 11-12 11-12 11-12 11-12 | 9.4 7.9 6.4 13.0 | 75.3 (70-79) 60.5 (55-68) 45.8 (40-52) 104.5 (84-123) | Spatial span Impaired MIQ & mild ID Preserved BIQ |
| Numminen et al. (2001) | NSID (n = 24) TDMA (n = 24) | 38.58 (49.88) 3-6 (5.50) | 5.63 5.67 | Raven | Corsi block Impaired |
| Vicari et al. (1995) | NSID (n = 14) TDMA (n = 24) | 16.7 5.7 | 51 5.5 | 39 101 | Forward visuo-spatial task – Corsi blocks Preserved DS & ID |
| Central Executive Visuospatial | | | | | |
| Alloway (2010) | BID (n = 39) | 7.11-11.7 <i>M</i> = 9.8 | | 70-85 <i>M</i> = 77 | WM (AWMA Alloway, et al., 2007) |
| | TDCA (n = 39) | 8.1-11.11 | | 99-133 <i>M</i> = 118 WASI | Spatial span Odd-one-out Impaired Impaired Impaired |
| Henry and Winfield (2010) | MID (n = 35) TDMA (n = 35) | 12-13 (12.6) 6-8 (7.5) | 7.7 7.8 | 57 101 | Odd one out span Preserved in same mental age Associated with number skills |
| Leken et al. (2011) | Mild ID (n = 10) TD (n = 5) | 275 27 | | 85-11 50-70 | Picture spam More complex Photos Drawing Objects Faces Impaired Impaired Impaired Impaired Impaired Impaired |
| Van der Molen et al. (2010) | Mild ID (n = 39) TDCA (n = 39) TDMA (n = 26) | 13-17 13-16 9-12 | 118.1 182.6 120.7 | 65 (55-75) (FIQ) 99.8 97.1 | Odd-One-Out (Henry 2001) Preserved MA Impaired CA # |
| Van der Molen et al. (2009) | Mild ID (n = 49) TDCA (n = 39) TDMA (n = 29) | 13-17 13-16 8-12 | | YIQ PIQ 68.6 71.9 98.6 98.5 100.5 97.7 | Out-One-Out Impaired CA Preserved MA |
| Henry and MacLean (2003) | NSID (n = 53) TDCA (n = 45) TDMA (n = 41) | 11-12 (11.11) 11-12 (12.1) 7-8 (7.11) | | 57.2 (40-79) 103.8 (84-123) 101.0 (82-136) | Odd-one-out-span Preserved MA |
| Henry and MacLean (2002) | NSID (n = 53) TDCA (n = 45) TDMA (n = 41) | 11-12 11-12 7-8 | | 57.2 103.8 101.0 | Odd one out Preserved MA impaired CA |

Table 1 (*Continued*)

| Henry (2001) | | BID (n = 10) | 11-12 | 9.4 | 75.3 (70-79) | Odd one out | Impaired |
|-------------------------------|--|------------------|-----------------|-------|-------------------------------|---|---|
| Vicari et al. (1995) | | Mild ID (n = 25) | 11-12 | 7.9 | 60.5 (55-68) | | |
| | | MI D (n = 21) | 11-12 | 6.4 | 45.8 (40-52) | | |
| | | TD CA (n = 22) | 11-12 | 13.0 | 104.5 (84-123) | | |
| Alloway (2010) | | NSID (n = 14) | 16.7 | 51 | 39 | | |
| | | TDMA (n = 24) | 5.7 | 5.5 | 101 | Backward visuo-spatial task | Preserved ID=TD |
| Phonological Executive | | | | | | | |
| Schuchardt et al. (2010) | | BID (n = 39) | 7.11-11.7 (9.8) | | 70-85 M = 77 | Backwards digit recall | Impaired |
| | | TDCA (n = 39) | 8.1-11.11 | | 99-133 M = 118 WASI | Listening recall Counting recall | Impaired Impaired |
| | | BID (n = 15) | 15 | | 82.63 (70-84) | Backward digit span | Impaired CA-preserved MA |
| | | MID (n = 15) | 15 | | 61.82 (50-69) | Counting span task | Impaired CA |
| | | NSTD (n = 25) | 15 | | 109.80 (90-115) | | Preserved MA deficits increased with the degree of ID |
| | | BID (n = 19) | 10 | | 81.21 | | |
| | | TDMA (n = 22) | 7 | | 104.91 | | |
| Van der Molen et al. (2010) | | Mild ID (n = 39) | 13-17 | 118.1 | 65 (55-75) (FIQ) | Backward digit recall | Preserved MA |
| | | TDCA (n = 39) | 13-16 | 182.6 | 99.8 | | Impaired CA |
| | | TDMA (n = 26) | 9-12 | 120.7 | 97.1 | Listening recall | Impaired |
| Maehler & Schuchardt (2009) | | NSID (n = 27) | 12.45 | | 75.44 | Backward digit spans | impaired |
| | | LD (n = 27) | 12.35 | | 100.08 | | ID=LD<TD impaired |
| | | TD (n = 27) | 10.58 | | 101.04 | Backwards word span Counting span | ID=LD-TD impaired ID=LD<TD |
| Van der Molen et al. (2009) | | Mild ID (n = 49) | 13-17 | | VIQ 68.6 PIQ 71.9 | Backward digit span | Impaired CA & MA |
| | | TDCA (n = 39) | 13-16 | | 98.6 98.5 | | Impaired Ca & MA |
| | | TDMA (n = 29) | 8-12 | | 100.5 97.7 | | |
| Numminen et al. (2001) | | NSID (n = 24) | 38-58 (49.88) | 5.63 | Raven | Digit span backwards | preserved |
| | | TDMA (n = 24) | 3-6 (5.50) | 5.67 | | | |
| Conners et al. (1998) | | NSID (n = 30) | 11.72 | 7.45 | 66.58 WISC-R verbal | Backward digit span | Impaired CA |
| | | TDCA (n = 30) | 12.36 | 7.45 | 107.42 | | Preserved MA |
| Vicari et al. (1995) | | TDVMA (n = 26) | 7.45 | | 106.77 | | CA>ID=MA |
| Numminen et al. (2002) | | DS (n = 15) | 16.6 | 5.3 | 36 | Backward digit span | Preserved ID=TD |
| Carreti et al. (2010) | | NSID (n = 14) | 16.7 | 51 | 39 | | |
| | | TDMA (n = 24) | 5.7 | 5.5 | 101 | | |
| Henry and Winfield (2010) | | NSID (n = 24) | 49.88 | | Raven | Digit Span Backward Task | Preserved |
| | | TDMA (n = 24) | 5.650 (3-6) | | | Verbal WM tests (Belachchi et al., 2010) | |
| | | NSID (n = 28) | 38.4 | 6.2 | Raven: 17.36 (IQ = 40-75) | | Impaired |
| | | TDMA (n = 28) | 6.5 | 6.6 | 17.50 | Backward word span Selective word span | Impaired |
| Henry and MacLean (2003) | | MID (n = 35) | 12-13 (12.6) | 7.7 | 57 | Listening span | Preserved in same mental age |
| | | TDMA (n = 35) | 6-8 (7.5) | 7.8 | 101 | | Associated with number skills |
| Henry and MacLean (2002) | | NSID (n = 53) | 11-12 (11.11) | | 57.2 (40-79) | Listening span | Preserved MA |
| | | TDCA (n = 45) | 11-12 (12.1) | | 103.8 (84-123) | Reverse digit span | Preserved MA |
| | | TDMA (n = 41) | 7-8 (7.11) | | 101.0 (82-136) | | |
| Henry (2001) | | NSID (n = 25) | 11-12 | | 57.2 | Listening span | Preserved MA |
| | | TDCA (n = 45) | 11-12 | | 103.8 | Reverse digit | impaired CA |
| | | TDMA (n = 41) | 7-8 | | 101.0 | | Preserved MA |
| Lanfranchi et al. (2002) | | BID (n = 10) | 11-12 | 9.4 | 75.3 (70-79) BAS 60.5 (55-68) | Listening span | impaired CA |
| | | Mild ID (n = 25) | 11-12 | 7.9 | 45.8 (40-52) | | Impaired |
| | | MI D (n = 21) | 11-12 | 6.4 | 104.5 (84-123) | Reverse digit span | Impaired |
| | | TD CA (n = 22) | 11-12 | 13.0 | | | |
| Danielson et al. (2013) | | NSID (n = 13) | 8-16 | 5.6 | LO | Backward word span | Impaired |
| | | TDMA (n = 30) | 5.2 | 5.6 | | Selective word recall | NSID<TD Impaired |
| | | NSID (n = 22) | 7.3 | 8 | | | EX. Verbal preserved |
| | | TDMA (n = 22) | 12 | | | | |
| | | TDCA (n = 22) | 14 | | | | |

Table 1 (Continued)

| | | | | | | |
|--------------------------------|------------------------|--|------|-------------------------------------|---|---|
| Carreti et al. (2010) | NSID (<i>n</i> = 28) | 38.4 | 6.2 | Raven: 17.36 (IQ = 40–75) | Dual span task | Impaired |
| | TDMA (<i>n</i> = 28) | 6.5 | 6.6 | 17.50 | | |
| Danielson et al. (2010) | NSID (<i>n</i> = 46) | 63.2 | | 62.8 Wechsler 100 | Dual task word recall: | Impaired |
| | TDCA (<i>n</i> = 92) | 63.2 | | | | |
| Maehtler and Schuchardt (2009) | NSID (<i>n</i> = 27) | 12.45 | | 75.44 | double span | All impaired |
| | LD (<i>n</i> = 27) | 12.35 | | 100.08 | | ID=LD<TD |
| | TD (<i>n</i> = 27) | 10.58 | | 101.04 | | |
| Van der Molen et al. (2007) | NSID (<i>n</i> = 50) | 15.3 | | Raven 55–85 | <u>Dual-task management</u> | |
| | TDCA (<i>n</i> = 25) | 15.3 | | | Dual task (Baddeley, 1997, verbal/visuospatial) | preserved CA and MA |
| | TDMA (<i>n</i> = 25) | 10.10 | | | | |
| Bayliss et al. (2005) | ID\LD (<i>n</i> = 50) | 13.6 (10.1–16.3) | 7.5 | 55.32 (27.72–69.89) | <u>Complex span tasks</u> Bayliss et al. (2005) | Preserved |
| | TDVMA (<i>n</i> = 50) | 7.3 (5.5–9.1) | 7.5 | BPVS-II 102.36 (80.81–129.17) | Verbal-verbal Verbal-visuo spatial | Preserved |
| Lanfranchi et al. (2002) | NSID (<i>n</i> = 13) | 8–16 | 5.6 | LO | Dual selective word recall | Impaired |
| | TDMA (<i>n</i> = 30) | 5.2 | 5.6 | | Dual visuospatial task | Impaired |
| | | | | | | DS=NSID < TDMA |
| Executive functions | | | | | | |
| Danielson et al. (2010) | NSID (<i>n</i> = 46) | 63.2 | | 62.8 Wechsler 100 | Tower of Hanoi | Preserved |
| | TDCA (<i>n</i> = 92) | 63.2 | | | Verbal fluency | Impaired |
| Van der Molen et al. (2007) | NSID (<i>n</i> = 50) | 15.3 | | Raven 55–85 | <u>Information retrieval and manipulation</u> | impaired CA |
| | TDCA (<i>n</i> = 25) | 15.3 | | | Word fluency test | Preserved MA |
| | TDMA (<i>n</i> = 25) | 10.10 | | | Planning WISC-R subtest | Impaired CA |
| | | | | | mazes Inhibition | Preserved MA |
| Conners et al. (1998) | NSID (<i>n</i> = 30) | 11.72 | 7.45 | 66.58 WISC-R verbal | digit generation (DG) Fluency | Impaired CA |
| | TDCA (<i>n</i> = 30) | 12.36 | 7.45 | 107.42 | | Preserved MA |
| | TDVMA (<i>n</i> = 26) | 7.45 | | 106.77 | | Impaired CA |
| Alloway (2010) | BID (<i>n</i> = 39) | 7.11–11.7 <i>M</i> = 9.8 | | 70.85 <i>M</i> = 77 | D-KEFS (Delly, 2001) | Preserved MA |
| | TDCA (<i>n</i> = 39) | 8.1–11.11 | | 99–133 <i>M</i> = 118 WASI | Shift attention Cognitive inhibition | Impaired |
| | | | | | Problem solving behavior | Impaired |
| Carreti et al. (2010) | NSID (<i>n</i> = 28) | 38.4 | 6.2 | Raven: 17.36 (IQ = 40–75) | Response inhibition | Impaired |
| | TDMA (<i>n</i> = 28) | 6.5 | 6.6 | 17.50 | <u>WM updating word span</u> (Palladino et al., 2001) | Impaired |
| Hartman et al. (2010) | B ID (<i>n</i> = 61) | 7–12 | | 71–79 | EF: | |
| | M ID (<i>n</i> = 36) | 7–12 | | 54–70 | Tower of London-TOL | Impaired |
| | TDCA (<i>n</i> = 97) | 7–12 | | 58.8 (50–67) | BADS-C Emslie 2003 | BID=MID<TDCA |
| Willner et al. (2010) | NSID (<i>n</i> = 20) | 39.9 | | BPVS: 81.6 | CEFA-ID Ball 2008 | Weighing up impaired |
| | TD (<i>n</i> = 10) | 38.6 | | 152.3 | | Decision making related more to EF than IQ |
| Bayliss et al. (2005) | ID\LD (<i>n</i> = 50) | 13.6 (10.1–16.3) <i>7.3 (5.5–9.1)</i> | 7.5 | 55.32 (27.72–69.89) | <u>Processing efficiency tasks (processing speed)</u> | |
| | TDVMA (<i>n</i> = 50) | | 7.5 | 102.36 (80.81–29.17) | Color association Object association TOH | ID/LD faster than TD Preserved ID more trials & violated more rules |
| Numminen et al. (2001) | NSID (<i>n</i> = 24) | 38–58 (49.88) | 5.63 | Raven | | |
| | TDMA (<i>n</i> = 24) | 3–6 (5.50) | 5.67 | | | |

NSID = ID with non-specific etiology, BID = borderline ID, Mild ID = mild ID with IQ 55–85, MID = moderate ID, TD = with typical development, CA = chronological age, MA = mental age TDMA = with typical development control matched by MA; TDCA = with typical development control matched by CA, VIQ = verbal IQ, PIQ = perceptual IQ, BAS = British Ability Scale, AWMA = Automated Working Memory assessment, D-KEFS = Delis-Kaplan Executive Function System, BADS-C = Children's version of the Behavioural Assessment of the Dysexecutive Syndrome, which includes six tests of EF, CEFA-ID = the Cambridge Executive Functioning Assessment for people with Intellectual Disability, WASI = Wechsler Abbreviated Scale of Intelligence

2.3. WM in individuals with NSID

2.3.1. Moderators related to background characteristics

2.3.1.1. CA versus MA as criteria for matching the ID and TD groups. The discrepancy between studies may originate in the criteria used for matching the groups. When matching between participants with ID and TD was by CA (*n* = 21), WM was impaired in all studies beyond modality and the component. Participants with TD exhibit a higher MA or IQ than their peers with ID, and it is not surprising that they outperform them on memory tasks. However, when matching between groups was based on MA (*n* = 41), 34% of the experiments yielded equal performance. To find what might explain the inferior

Table 2

The hierarchy of WM tasks among participants with NSID.

| Process type | Preserved | Task | N | N Preserved |
|----------------------------------|-----------|--------------------|----|----------------|
| Phonological loop (n = 68) | 40% | Digit | 22 | 9 |
| | 20% | | | |
| | 71% | | | |
| | 77% | Words | 15 | 3 |
| | 13% | Similarity effect | 7 | 5 |
| | | Word length effect | 9 | 7 |
| | | Non-words | 15 | 2 |
| | | Visual | 15 | 7 |
| | | Spatial | 17 | 6 |
| | | Visuospatial tasks | 17 | 6 |
| visuo-spatial sketch pad (32) | 46% | Exec. Phonological | 24 | 9 |
| | 35% | Digit backward | | |
| | 22% | Words backward | 9 | 2 |
| | 50% | Word Fluency | 4 | 2 |
| | 60% | Planning | 5 | 3 |
| | 0 | Inhibition | 3 | 0 |
| | 0 | Switching verbal | 2 | 4 |
| | 40% | WM dual task | 10 | 3 |
| Central executive (n = 57) | | | | |

performance in the remaining experiments, we triangulated between the studies and found other moderators that influence memory performance of individuals with ID, even when the groups are matched by MA.

Furthermore, memory was preserved in accordance to the intelligence tests used for matching between the groups across all modalities and components of Baddeley's model (phonological loop, visuospatial and executive function tasks). When the criteria for matching groups was passive vocabulary (Peabody Picture Vocabulary- PPVT; Dunn & Dunn, 1981; Dunn, Dunn, Whetton, & Burley, 1997; Wechsler Intelligence Scale for Children, WISC-R-III; Wechsler, 1991; British Ability Scale; Elliott, Smith, & McCulloch, 1996), the performance of participants with ID was inferior to that of participants with TD. However, when matching was based on fluid intelligence tests (Standard progressive matrices, Raven, Court, & Raven, 1992), logical thinking (Columbia Mental Maturity Scale; Burgemeister, Hollander, & Lorge, 1972), the Logical Operation Test (Vianello & Marin, 1997) or Leiter Performance Intelligence (Roid & Miller, 1997), performance was preserved (see Bayliss et al., 2005; Van der Molen et al., 2010).

2.3.1.2. Task-related moderators – phonological loop. Of 68 phonological experiment tasks, 27% were preserved (including digit, words and non-words tasks). We found the following hierarchy in the phonological loop from more to less preserved): Digit span, memory of words and memory of non-words.

2.3.1.3. Digit span. In 40% of 22 experiments, the performance of participants with NSID was preserved only when compared to individuals with TD with the same MA. Furthermore, the discrepancy between the preserved and impaired experiments stems, among others, from the intelligence tests used for matching between the groups, the level of ID and the scoring methods. When the criteria for matching groups was passive vocabulary (i.e., Peabody Picture Vocabulary-PPVT; Dunn & Dunn, 1981), the performance of participants with ID was inferior to that of participants with TD. However, when matching was based on fluid intelligence tests (Standard progressive matrices test; Raven et al., 1992) or logical thinking (Columbia Mental Maturity Scale; Burgemeister et al., 1972), memory was preserved. Furthermore, in Bayliss et al. (2005), for example, digit span scores of participants with ID was lower than among participants with TD. This study included participants with both mild and moderate ID. In Henry and Winfield (2010), who found that participants' performance for digit memory was impaired, scoring was based on the number of successful attempts needed to complete the task, which Ferguson, Bowey, and Tilley (2002) consider more reliable. When scoring was based on the longest list participants were able to recall (Henry & MacLean, 2003), performance was preserved.

2.3.1.4. Words. Phonological memory measured by word span was preserved in 20% of 15 experiments when matching between the groups was based on fluid intelligence (Carretti, Belacchi, & Cornoldi, 2010; Hasselhorn & Mähler, 2007; Lanfranchi, Cornoldi, & Vianello, 2002), and impaired when matching was based on verbal tests. These criteria were true also in word length and similarities effects. Word length tests examine the effects of word length (Service, 1998) on phonological memory span and spontaneous sub-vocal rehearsal ability. In 77% (n = 9) of the experiments, participants with ID were able to recall words with 1 or 2 syllables, but performance for longer words was not preserved. The *similar/dissimilar words effect* was preserved in 71% of 7 experiments. This finding indicates a severe deficit in the phonological store that is solely attributable to storage capacity rather than processing accuracy, which Schuchardt et al. (2011) consider as a core deficit in the population with ID (see Discussion Section).

2.3.1.5. Non-words. *Non-words* (repetition) was preserved in only 13% of 15 experiments, in two studies in which matching between the groups was according to MA and when a fluid intelligence test (Standard progressive matrices, [Raven et al., 1992](#)) was used for matching between the groups.

2.4. Visuospatial sketchpad

2.4.1. Moderators related to background characteristics

2.4.1.1. CA versus MA as criteria for matching the ID and TD groups. When matching was according to MA, the performance of participants with ID was equal to the controls with TD in 40% of 32 experiments. In the preserved experiments, fluid intelligence tests (i.e., Standard progressive matrices) were used for matching between the groups.

2.4.1.2. Task-related moderators. Memory of visual-object was measured ([Visu-Petra & Benga, 2007](#)) by visual similarity, visual complexity, and span/visual patterns tests. The findings indicate that visual object was preserved in 43% of 15 experiments ([Table 2](#)). Memory was impaired when comparison was based on CA, but was more preserved when comparison was based on MA. Pattern span was preserved. However, matrixes revealed impairment. Spatial performance was measured with the memory span for locations, Corsi block simple and complex tasks (where the distance between blocks is larger than in simple tasks). Spatial memory was preserved in 35% of 17 experiments ([Table 2](#)). Most of the experiments used the Corsi block ([Millner, 1971](#)). Memory was impaired when comparison was based on CA, but more preserved when comparison was based on MA. Corsi complex was impaired.

2.5. Central executive

2.5.1. Moderators related to background characteristics

2.5.1.1. CA versus MA as criteria for matching the ID and TD groups.. In 2 of the 57 experiments, memory was preserved when matching was according to CA. [Van der Molen et al. \(2007\)](#) examined a visual-verbal dual task among adolescents with mild ID versus participants with TD with the same CA and participants with TD with the same cognitive level based on the Raven's Standard Progressive Matrices Test ([Raven et al., 1992](#)). However, this experiment was conducted in two stages: The phonological and visuospatial tasks were administered separately, after which they were administered as a dual task. It is possible that the previous familiarity with the task contributed to the preserved performance of the participants with ID on the dual task. [Danielsson et al. \(2010\)](#) examined central executive performance among adults with NSID versus adults with TD with the same CA. Tower of Hanoi, word fluency, dual task and word recall were administered. Participants with ID exhibited preserved performance only for Tower of Hanoi. According to Miyake et al. (2000), this task requires planning, which is considered to have a lower level of task load than updating or shifting tasks. When matching was according to MA, the executive performance of participants with NSID was preserved in 72% of 35 experiments.

2.5.1.2. Task-related moderators. We counted the number of preserved/impaired experiments in central executive tasks and found the following hierarchy (from more to less preserved): Visuospatial tasks followed by phonological tasks and dual tasks. In 35% of the visuospatial executive tasks the performance of participants with NSID was preserved. In the *phonological executive tasks*: Digit memory was preserved in 37% of 24 experiments, indicating that remembering a digit is easier for the population with ID. Memory of executive words tasks was preserved in 22% of the nine experiments that examined executive phonological tasks. In [Henry and MacLean \(2002, 2003\)](#) and [Henry and Winfield \(2010\)](#), participants were asked to repeat the last word of sentences, without reference to their order. However, in [Van der Molen et al. \(2009, 2010\)](#), they were asked to repeat the last word of each sentence in the order they were presented, which increases the level of task control and therefore memory was impaired. The findings show that participants with ID exhibit greater difficulty with listening span, selective word span and backward word span tests, which require simultaneous storage and processing of phonological information, than participants with TD with the same MA ([Carretti et al., 2010; Lanfranchi et al., 2002](#)).

2.5.1.3. Dual task. Dual tasks involve memory performance of two modalities simultaneously (phonological-phonological, spatial-spatial or phonological-spatial), and require a higher level of control ([Baddeley, 2007](#)). Performance was preserved in 44% of nine experiments for the verbal-verbal and verbal-visual-spatial dual tasks ([Bayliss et al., 2005; Van der Molen et al., 2007](#)). In [Carretti et al. \(2010\)](#) and [Lanfranchi et al. \(2002\)](#), the secondary tasks were performed only in response to a stimulus, requiring the participant to focus attention on two sources simultaneously, and performance was impaired.

2.5.1.4. Other executive functions. Preserved performance was found for planning (Tower of Hanoi, WISC-R subtest mazes), although the number of errors and moves was much higher among participants with NSID than among participants with TD. Performance was also preserved in retrieval, verbal fluency and processing efficiency tasks. Impaired performance was found in updating word span tasks ([Carretti et al., 2010](#)) that require focusing attention simultaneously on two actions ([Palladino, Cornoldi, De Beni, & Pazzaaglia, 2001](#)), inhibition and switching. None of these tasks were preserved. These tasks demand not only storage, but also a high level of control, constant retrieval of information and other manipulations beyond memory itself.

In conclusion, we found a hierarchy of WM performance in the population with NSID (from more to less preserved): visuospatial sketchpad, phonological loop and some of the central executive functions (planning, fluency and even dual tasks), with phonological loop tasks found at the bottom of the hierarchy.

3. Discussion

The intriguing message to emerge from this integrative review is that WM performance of individuals with ID is context-sensitive (Roediger, 2008). That is, there are some conditions in which WM is relatively more preserved and some conditions in which it is impaired. Different abilities are preserved to various degrees. This is the first message of this review.

Two fundamental questions will be at the core of the discussion: (a) What are the moderators that govern the preservation or impairment of WM performance among participants with NSID? (b) Whether and what are the common denominators between the processes of long-term memory (LTM) and WM in the population with NSID?

Our review of WM studies in the population with NSID revealed similarities and differences to the findings that emerged from our long-term explicit memory studies review (Lifshitz-Vahav & Vakil, 2014). Our review suggests a taxonomy of two clusters of moderators that influence WM performance in the population with NSID: (a) Participant-related moderators (CA, intelligence level and etiology), and (b) Task-related moderators.

The participants cluster includes the same moderators or clusters that govern long-term explicit memory in the population with ID: CA and MA (or intelligence). However, the participant-related moderators are quite different due to the specific construct of the WM. Our findings support the horizontal and vertical model of WM (Cornoldi & Vecchi, 2003) in the population with NSID. That is, WM performance in this population is influenced by the modality according to the components model (phonological loop, the visuospatial sketchpad and the central executive which is involved in conditions of high level processing of the two other slave systems) of Baddeley et al. (2011), but is also influenced by the cognitive load of the tasks in each component. Our claim is that only the interrelation between the two moderators provides an adequate profile of WM in individuals with ID. This is the second message of our research review. We will now relate to the two clusters of moderators.

3.1. Cluster 1: participant-related moderators

3.1.1. The influence of CA on WM in the population with ID

Similarly to the moderator that governs explicit memory in the population with ID, this review of WM studies indicates that CA plays an important role in determining WM performance of individuals with NSID. Although the influence of CA on WM in the population with ID was not examined directly in the WM studies (most of the experiments focus on adolescents with NSID), its role emerges from the difference in performance when selection of the control group was based on MA rather than on CA. In most of the WM studies that matched individuals with ID and controls by CA, individuals with TD outperformed their peers with ID. This is self-evident. Thus, the different gap pattern between the two groups when selection of the latter was based on CA implies the effect of age on WM.

However, in some of the studies that matched participants with ID and controls with TD based on MA, the performance of participants in the three etiologies was equal to their controls. When matching is based on MA, the participants with ID are older than their peers with TD, which gives them an advantage (Van der Molen et al., 2010). This finding raises the question of a developmental-difference controversy (Zigler & Balla, 1982) in the population with ID. The question is whether the deficit in WM can be interpreted as reflecting a developmental lag or a qualitative deviation (structural difference) from typical development (Zigler & Balla, 1982). The fact that the performance of individuals with NSID is equal to that of their controls with TD when the former are older supports the developmental approach. Carlesimo, Marotta and Vicari (1997) and Facon and Facon-Bollengier, (1999) found that adolescents with ID outperformed children with ID with the same MA in crystallized tests. Lifshitz and Katz (2009) and Lifshitz, Tzuriel, Weiss, and Tzemach (2010) found a similar trend among adults and adolescents with ID with the same cognitive level. The Compensation Age Theory (Lifshitz-Vahav, 2015) postulates that CA determines the cognitive ability of individuals with ID beyond their MA. Their maturity and cumulative life experience may explain their more efficient coping with cognitive skills in adulthood (Numminen, Lehto, & Ruoppila, 2001).

3.1.2. The influence of MA or intelligence on WM in the population with ID

Jarrold, Baddeley and Phillips (2007) stated that IQ may be more important in predicting performance on LTM tasks, which are considered a fluid ability and are more related to general intelligence *g* than to verbal MA. Our review of WM studies in the population with ID supports this claim: In all experiments, when memory performance was inferior to that of participants with TD, matching was based on verbal ability or the MA (examined by the PPVT, Dunn & Dunn, 1981, 1997). In all experiments that indicate equal performance between participants with NSID and TD, matching was based on fluid intelligence tests (examined by the Standard progressive matrices, Raven et al., 1992; or the Logical Operations, Vianello & Marin, 1997). This discrepancy in WM performance as a consequence of the different tests used for matching between participants with ID and TD was also supported in our previous explicit memory studies review (Lifshitz-Vahav & Vakil, 2014). Baddeley et al. (2011) claim that the three components of WM are associated with fluid intelligence. Thus, it is not surprising that fluid intelligence tests are more accurate and reliable measures of the basic cognitive intelligence of individuals with ID, and that they predict both their WM and LTM performance better than their MA.

Our findings support the vertical model suggested by [Cornoldi and Vecchi \(2003\)](#). This model reflects the degree of required control, which is defined by the amount of active processing required to manipulate the information maintained in a temporary memory system. The vertical model of cognitive load raised by [Cornoldi and Vecchi \(2003\)](#) is parallel to the concept of attention level in the time-based resources-sharing model suggested by Camos and Barrouillet (2014, see p. 4). This WM integrative review indicates that individuals with ID can manage with WM tasks demanding a lower level of attention in the process and storage stages. However, they exhibit difficulties in highly attention-demanding executive tasks involving other manipulations besides the recall itself, such as selection, retrieval, updating, or inhibition. Our findings indicate a taxonomy of task performance in each component in the model by [Baddeley et al. \(2011\)](#). We identified three levels of cognitive load or task control in the phonological loop, the visuospatial sketchpad and the executive function. Consistent with our hypothesis, memory performance was equal to the performance of participants with TD at a lower cognitive load or lower attention-demanding tasks, but was inferior to participants with TD in higher cognitive load tasks which require a higher level of attention.

The fact that the performance of individuals with NSID depends on the task load indicates that WM performance does not depend solely on the modality, but rather on the participants' basic cognitive level, that is – on their mental age, but more correctly on their intelligence (considering their chronological age). This is the third message of our study. This claim is in accordance with that of [Levy \(2011\)](#) and [Levy and Antebi \(2004\)](#) who found that IQ was the best predictor of word reading skills among participants with NSID, Down syndrome and Williams syndrome, beyond the unique phenotype of each etiology. We shall now present the hierarchy of the three components in the population with ID.

3.2. Cluster 2 – task-related moderators

The findings indicate a hierarchy (from more to less preserved) in performance of individuals with ID of three components: The visuospatial tasks, some of the executive functions tasks (it should be noted that as in qualitative studies, this hierarchy is based on percent and not on sample and effect size as used in meta-analyses, see Procedure Section), and the phonological loop tasks at the bottom of the hierarchy (least preserved).

3.2.1. The visual component

The findings of the more preserved WM visual tasks were also supported in our integrative research review of explicit LTM studies in the population with ID ([Lifshitz-Vahav & Vakil, 2014](#)). In the LTM studies, none of the verbal tasks yielded equal performance between individuals with ID and their TD controls. Equal performance on verbal tasks was achieved when the word-list was accompanied by visual scaffolding. The gap between memory of visual and verbal memory in WM as well as in LTM supports the 'picture superiority' hypothesis (Dulaney & Ellis, 1991; Paivio, 1971) claiming that processing visual information enables individuals with ID to develop the meaning behind the visual stimuli more than words alone. Visual pattern span was more preserved than both simple and complex matrixes, that were impaired.

3.2.2. Spatial memory

Visual spatial WM memory was less preserved compared to visual object tasks even when ID and TD groups were matched by MA. The essence of spatial memory has occupied scientific literature since the 1970s. Hasher and Zacks (1979) claimed that spatial memory is an automatic process. They presented six criteria for automatic processing and claimed that spatial memory is not influenced by different moderators related to the task or to the subjects (task demands, practice, feedback, awareness of intent, CA and intelligence level). In contradistinction, Naveh-Benjamin's (1987, 1988) findings indicate that encoding of spatial location information is influenced by these moderators, and this memory therefore involves effort.

Based on our long-term explicit memory review ([Lifshitz-Vahav & Vakil, 2014](#)) and the present WM review, our argument is that encoding of location memory in individuals with NSID is mediated by non-automatic processes (Naveh-Benjamin, 1988) and is influenced by task load (Vinter & Detable, 2003), strategy use and intentional learning (Alevriadou, 2010; Dulaney & Ellis, 1991). According to Dulaney, Raz and Devine (1996) and Alevriadou (2010), encoding of spatial information is mediated by effort processes, and spatial memory is processed differently among people with ID than among those with TD. The population with ID relies on minimal cognitive resources, and what seems to be an automatic process for those with TD who have sufficient cognitive resources necessitates an effort among those with ID. This conclusion should be regarded with caution, due to the small number of studies on this issue.

3.2.3. Phonological loop

Similarly to our integrative research review of long-term memory studies in the population with ID ([Lifshitz-Vahav & Vakil, 2014](#)), this integrative research review of WM studies in the population with ID indicates that the phonological verbal tasks are at the bottom of the performance hierarchy compared to the visuospatial sketchpad and some of the executive function tasks. It should be noted that in our long-term explicit memory integrative research review ([Lifshitz-Vahav & Vakil, 2014](#)), verbal tasks yielded equal performance of individuals with ID and TD only with visual scaffolding. The findings of this WM review support our previous claim about visual images superiority in the population with NSID. Unlike our integrative research review of explicit LTM, the current integrative WM review indicated that of 68 phonological experiment tasks, 27% were preserved (including digit, words and non-words tasks). That is, in the short term, individuals with NSID could hold verbal material without visual scaffolding.

Task hierarchy in the phonological loop was as follows (from more to less preserved): Digit span forward, followed by words and then non-words. The relatively preserved memory for a digit can be explained by Piaget's (1970) cognitive development stages. Memory for numbers is acquired at the relatively young age of three, prior to memory for words. The first ten single-digit numbers are preserved. Recalling numbers is accompanied by rhythmic movement, which facilitates the remembering of numbers, also for individuals with ID (Vygotsky, 1986). Another explanation could be the feature of the phonological loop which is the capacity for vocal rehearsal. In this system, sequences of well-learned items such as a digit are readily retrievable from the LTM and may be maintained before their constituent features are disrupted by decay or interference. Thus, as stated, contrary to the findings of the long-term memory review (Lifshitz & Vakil, 2014), individuals with NSID could relatively hold verbal material in the STM without visual scaffolding (Schuchardt et al., 2011).

Our study revealed the word length effect, i.e. short words take a shorter time to articulate and benefit from more rehearsals, which increases the probability that they will be recalled (Jarrold & Hall, 2013; Service, 1998). Our findings indicate that individuals with ID are capable of recalling two-syllable words, and in some cases three syllables, when other moderators are employed. Words with four syllables place more constraints on memory storage, and decrease the performance, even of participants with TD (Schuchardt et al., 2011).

According to Hasselhorn and Mähler (2007) and Schuchardt et al. (2011), this finding supports the claim of automatic activation of the sub-vocal rehearsal. That is, individuals with ID show no specific deficits in the rehearsal process. However, our findings reveal a deficit in repetition of non-words (familiarity or meaningfulness effect) of different syllable lengths. Lack of familiarity or meaning causes a barrier on the storage mechanism compared to the population with TD (Schuchardt et al., 2011). This finding indicates a severe deficit of individuals with ID in the phonological store which can be attributed to storage capacity and not to the accuracy of processing (Schuchardt et al., 2011). It is documented that storage plays a key role in early language development, in the acquisition of active vocabulary, and in the use of grammatical forms and syntactic structures (Adams, 1996; Baddeley, Gathercole, & Papagno, 1998). Limited storage capacity encumbers vocabulary and grammatical structures, which are the core constraint in the cognitive development of individuals with ID. When the expansion rate of the knowledge base is slow, less content is entered, processed, and subsequently stored in long-term and working memory. According to Schuchardt et al. (2011), this indicates a qualitative difference between individuals with ID compared to those with TD. On the other hand, knowledge of the lexicon and of grammatical structures can compensate for incomplete word representations in the phonological store (Brown, Preece, & Hulme, 2000).

Individuals with NSID exhibit the phonological similarity effect, i.e. storage of similar words to more confusion than dissimilar words (Camos & Barrouillet, 2014). Concurrent articulation using similar language processes could block or at least impede the sub-vocal rehearsal of memory items and impair WM recall (Camos & Barrouillet, 2014).

This finding indicates accuracy of processing in the population with ID (Hasselhorn & Mähler, 2007; Jarrold, Baddeley, & Hewes, 2000; Rosenquist, Conners, & Roskos-Ewoldsen, 2003).

In conclusion, our review revealed that the deficit of individuals with NSID does not lie in rehearsal or accuracy of processing of verbal material but in the phonological store, which can be attributed to storage capacity. In this respect, the deficit in the phonological loop of individuals with NSID is structural (Schuchardt et al., 2011). Deficits in storage capacity can therefore be seen as a causal factor in intellectual disabilities (Schuchardt et al., 2011). Limited storage capacity causes difficulty in the acquisition of vocabulary and grammatical structures. The language problems that people with ID typically experience, which seem to have a negative impact on general cognitive development, can be seen in this context. Consequently, the expansion rate of the knowledge base is slow – less content is entered, processed, and subsequently stored in LTM. It thus seems likely that both a lack of phonological storage capacity and the associated linguistic deficits contribute to the cognitive deficits observed in individuals with mild NSID. More studies are needed in order to support this claim.

3.2.4. Executive function

The same explanations that we proposed earlier for the more preserved digit span forward and words can serve as an explanation for their relatively more preserved digit backward, letter and word fluency. These abilities are readily retrievable from LTM and are less vulnerable to decay. Participants with NSID also exhibit more preserved performance in planning tasks (verbal or visuospatial), which are considered low-demand tasks, even though they are classified as executive function tasks.

Thus, participants with ID are capable of coping with executive function tasks, albeit at a lower control level beyond the modality. However, they exhibit lower performance compared to participants with TD in higher control/attention tasks. For example: Selecting requires making a semantic judgment and remembering (Van der Molen et al., 2009, 2010) or selecting an item according to a specified criterion and remembering (Palladino, 2001). Updating requires removing some or all of the items stored in WM and replacing them with new items. Shifting requires refreshing tasks in which a cue directs the subject to foreground a recently presented item. Inhibition is the ability to ignore irrelevant stimuli while performing an additional task. It includes the ability to change and/or stop an action, behavior or response when this is not desirable. Dual tasks require relating to two modalities simultaneously. Although these tasks differ in their nature, they are well-known for capturing a higher attention level because they require cognitive manipulation beyond the memory itself (Camos & Barrouillet, 2014).

The role of rehearsal in WM is controversial. Earlier we presented the approach that associated sub-vocal rehearsal with WM. However, Jarrold and Hall (2013) state that rehearsal plays a role in LTM. That is, it is needed for assisting people to recall after a delay. As stated by Jarrold and Hall (2013): "Rehearsal is not required when one recalls a list of just-presented items immediately... However, in situations in which individuals have to maintain the information during an interval, rehearsal

could be employed. . . An individual's rehearsal capacity is constrained by, and may in fact be secondary to, his or her recall capacity" (p. 184). In our opinion, a meta-analysis that will compare between WM studies in the population with NSID and LTM in this population should be conducted in order to shed light and give a more precise answer to this issue.

4. Conclusions, limitations, and future research

Our review indicates similarities and differences between WM and LTM (Lifshitz-Vahav & Vakil, 2014) performance mechanisms in the population with ID. Two clusters of moderators influence the performance of the two types of memory: The participants cluster includes CA and intelligence. However, the participant-related moderators are quite different due to the specific construct of the WM. WM performance in the population with NSID is dictated by the component (Baddeley & Hitch, 1974; Baddeley et al., 2011) or modality. The visuospatial sketchpad was more preserved, followed by several executive tasks. The phonological loop was at the bottom of the hierarchy. However, WM performance is governed not only by modality, but also by the cognitive load of the task (Lanfranchi, Carretti, Spanò, & Cornoldi, 2009; Lanfranchi, Cornoldi, Drigo, & Vianello, 2009; Vertical-Horizontal Model, Cornoldi & Vecchi, 2003). This finding indicates that intelligence plays an important role in determining WM performance of individuals with ID beyond modality. We found a hierarchy in task performance in each component (from more to less preserved): In the phonological loop: digit span, followed by memory of words and memory of non-words. In the visuospatial sketchpad: memory of visual tasks and then memory of spatial tasks. In the central executive: Perceptual planning and retrieval (fluency) and some of the dual tasks were more preserved than shifting, inhibition and selection. The executive phonological tasks were at the bottom of the performance hierarchy.

One factor possibly limiting the current analysis is that we relied on journal articles. Additional non-significant results were perhaps not included in published articles or dissertations due to the 'file drawer problem' which is problematic in all research, and in reviews in particular. The effect of CA on WM was examined with reference to the control group. However, differences in WM of participants with ID and controls with TD of various ages could not be examined due to insufficient memory studies among adults (and children) with NSID. Using different types of intelligence tests for matching (visual/verbal) between ID and TD groups may yield different findings. We recommend that future researchers use more than one intelligence test for matching between the groups. Conducting integrated research that will examine WM and long-term memory will help solve the source of deficit in verbal tasks in the population with NSID. This integrative review focused on individuals with NSID. There is a need for more research on Williams syndrome, Down syndrome and other etiologies. Research examining the effect of gender on memory is recommended (Perez, Peynircioğlu, & Blaxton, 1998). The hierarchy in WM performance in the three components was based on counting the number of preserved/impaired experiments in each buffer. It is recommended to conduct a meta-analysis of LTM and WM which will take the sample and effect size of the studies in both types of memory into account. A mathematical-analytical method that will define the cognitive load of the various tasks in a more systematic way is needed.

4.1. Educational implications

Educators should be aware of the notion that CA affects memory and strategy use. They should keep in mind that the memory of participants with ID could improve with age. Therefore, in instances when students with NSID do not successfully learn material when it is first presented, it is possible that they will be able to master it at a later age. Such awareness should encourage teachers to continue training and practicing strategies with students despite the lack of immediate effects. Visual material is encoded more deeply and creates a visual image in addition to the symbolic meaning of the word (Paivio, 1971). Visual scaffolding is recommended when teaching individuals with ID. Rehearsal leads to better recall and deep processing (Atkinson & Shiffrin, 1968) and was found to be an efficient manipulation in the population with ID (Brock, Brown, & Boucher 2006). This leads to the following practical recommendations: When teaching prose, narratives, or current events, use Luftig's (1987) task analysis method: divide the narrative into small units or paragraphs. Repeat each unit several times. Encourage participants to rehearse the words or story they heard verbally. Employ repetitions. When teaching social problem solving, repeat the problem and the solution several times. When giving mathematical exercises, repeat each exercise several times (Luftig, 1987). The required task control should be taken into account when preparing intervention programs for these etiologies.

Acknowledgement

This work was supported by a grant from the Shalem Fund for the Development of Services for People with Intellectual Disabilities in the Local Councils in Israel.

References

- Alloway, T. P., Gathercole, S. E., & Pickering, S. J. (2007). *Automated working memory assessment*. London: Psychological Corporation.
- Alloway, T. P. (2010). Working memory and executive function profiles of individuals with borderline intellectual functioning. *Journal of Intellectual Disability Research*, 54, 448–456.
- Atkinson, R. C., & Shiffrin, R. M. (1968). Human memory: A proposed system and its control processes. In W. E. Spence, & J. T. Spence (Eds.), *The psychology of learning and motivation: Vol. 2. Advances in research and theory* (pp. 89–195). New York: Academic.

- Atkinson, R. C., & Shiffrin, R. M. (1971). The control of short-term memory. *Scientific American*, 225, 82–90.
- Baddeley, A. D., & Hitch, G. (1974). Working memory. *The Psychology of Learning and Motivation*, 8, 47–90.
- Baddeley, A. D., & Logie, R. H. (1999). Working memory: The multiple component model. In A. Miyake, & P. Shah (Eds.), *Models of working memory: Mechanisms of active maintenance and executive control* (pp. 28–61). Cambridge: Cambridge University Press.
- Baddeley, A. D., Allen, R. J., & Hitch, G. J. (2011). Binding in visual working memory: the role of the episodic buffer. *Neuropsychologia*, 49, 1393–1400.
- Baddeley, A. D., Thomson, N., & Buchanan, M. (1975). Word length and the structure of short-term memory. *Journal of Verbal Learning and Verbal Behaviour*, 14, 575–589.
- Baddeley, A. D., Gathercole, S., & Papagno, C. (1998). The phonological loop as a language learning device. *Psychological Review*, 105, 158–173.
- Baddeley, A. D. (1997). *Human memory: Theory and practice*. UK: Psychology Press.
- Baddeley, A. D. (2000). The episodic buffer: A new component of working memory? *Trends in Cognitive Science*, 4, 417–423.
- Baddeley, A. D. (2003). Working memory: Looking back and looking forward. *Neuroscience*, 4, 829–839.
- Baddeley, A. D. (2007). *Working memory, thought and action*. New York: Oxford University Press.
- Baddeley, A. D. (2008). What's new in working memory? *Psychology Review*, 13, 2–5.
- Bayliss, D. M., Jarrold, C., Baddeley, A. D., & Leigh, E. (2005). Differential constraints on the working memory and reading abilities of individuals with learning difficulties and typically developing children. *Journal of Experimental Child Psychology*, 92, 76–99.
- Belacchi, C., Carretti, B., & Cornoldi, C. (2010). The role of working memory and updating in Coloured Raven Matrices performance in typically developing children. *European Journal of Cognitive Psychology*, 22, 954–1080.
- Bellugi, U., Lichtenberger, L., Jones, W., Lai, Z., & St George, M. (2000). The neurocognitive profile of Williams syndrome: A complex pattern of strength and weaknesses. *Journal of Cognitive Neuroscience*, 12, 7–29.
- Borkowski, J. G., Carr, M., & Pressley, M. (1987). 'Spontaneous' strategy use: Perspectives from metacognitive theory. *Intelligence*, 11, 61–75.
- Brown, G. D. A., Preece, T., & Hulme, C. (2000). Oscillator-based memory for serial order. *Psychological Review*, 107, 127–181. <http://dx.doi.org/10.1037/0033-295X.107.1.127>
- Burgemeister, B. B., Hollander, L. H., & Lorge, I. (1972). *Columbia mental maturity scale* (3rd ed.). San Antonio, TX: Psychological Corporation.
- Camos, V., & Barrouillet, P. (2014). Attentional and non-attentional systems in the maintenance of verbal information in working memory: The executive and phonological loops. *Frontiers in Human Neuroscience*, 7 <http://dx.doi.org/10.3389/fnhum.2014.00900>
- Campione, J. C., & Brown, A. L. (1984). Learning ability and transfer propensity as sources of individual differences in intelligence. In P. H. Brooks, R. Speber, & C. McCauley (Eds.), *Learning and cognition in the mentally retarded* (pp. 265–293). Hillsdale, NJ: Erlbaum.
- Carlesimo, G. A., Marotta, L., & Vicari, S. (1997). Long-term memory in mental retardation: Evidence for a specific impairment in subjects with Down's syndrome. *Neuropsychologia*, 35, 71–79.
- Carretti, B., Belacchi, C., & Cornoldi, C. (2010). Difficulties in working memory updating in individuals with intellectual disability. *Journal of Intellectual Disability Research*, 54, 337–345. <http://dx.doi.org/10.1111/j.1365-2788.2010.01267.x>
- Conners, F. A., Carr, M. D., & Willis, S. (1998). Is the phonological loop responsible for intelligence-related differences in forward digit span. *American Journal on Mental Retardation*, 103, 1–11.
- Conway, A. R. A., Kane, M. J., Bunting, M. F., Hambrick, D. Z., Wilhelm, O., & Engle, R. W. (2005). Working memory span tasks: A methodological review and user's guide. *Psychonomic Bulletin and Review*, 12, 769–786.
- Cooper, H. M. (1982). Scientific guidelines for writing integrative research review. *Review of Education Research*, 52, 291–302.
- Cornoldi, C., & Vecchi, T. (2003). Visuo-spatial working memory and individual differences. Hove, UK: Psychology.
- Cornoldi, C., Rigoni, F., Venneri, A., & Vecchi, T. (2000). Passive and active processes in visuo-spatial memory: Double dissociation in developmental disabilities. *Brain and Cognition*, 43, 117–120.
- Danielsson, H., Henry, L., Rönneberg, J., & Nilsson, L. (2010). Executive functions in individuals with intellectual disability. *Research in Developmental Disabilities*, 31, 1299–1304.
- Danielsson, H., Henry, L., Messer, D., & Rönneberg, J. (2012). Strengths and weaknesses in executive functioning in children with intellectual disability. *Research in Developmental Disabilities*, 33, 600–607.
- Dunn, L. M., & Dunn, L. M. (1981). *Peabody pictures vocabulary test, revised (PPVT)*. Circle Pines, MN: American Guidance Service.
- Dunn, L. M., Dunn, L. M., Whetton, C., & Burley, J. (1997). *The British picture vocabulary scale* (2nd ed.). Windsor: NFER-Nelson..
- Elliott, C. D., Smith, P., & McCulloch, K. (1996). *British Ability Scales—BAS II* (2nd ed.). London: Nelson.
- Ellis, N. R. (1978). Do mentally retarded have memory deficits? *Intelligence*, 2, 41–45.
- Facon, B., & Facon-Bollengier, T. (1999). Chronological age and crystallized intelligence of people with intellectual disability. *Journal of Intellectual Disability Research*, 43, 489–496.
- Ferguson, A. N., Bowey, J. A., & Tilley, A. (2002). The association between auditory memory span and speech rate in children from kindergarten to sixth grade. *Journal of Experimental Child Psychology*, 81, 141–156.
- Fink, W., & Cegelka, P. T. (1982). Characteristics of the moderately and severely mentally retarded. In P. T. Cegelka, & H. Y. Prehm (Eds.), *Mental retardation* (pp. 231–259). Columbus, OH: Merrill.
- Garlick, D., & Sejnowski, T. (2006). There is more to fluid intelligence than working memory capacity and executive function. *Behavioral and Brain Sciences*, 29, 134–135. <http://dx.doi.org/10.1017/S0140525X0630903X>
- Gathercole, S., & Alloway, T. P. (2006). Practitioner review: Short-term and working memory impairments in neurodevelopmental disorders: Diagnosis and remedial support. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 47(1), 4–15.
- Gathercole, S. E., Adams, A.-M., & Hitch, G. J. (1994). Do young children rehearse? An individual-differences analysis. *Memory and Cognition*, 22, 201–207. <http://dx.doi.org/10.3758/BF03208891>
- Gathercole, S. E., Alloway, T. P., Willis, C., & Adams, A. M. (2004). Working memory in children with literacy and mathematical difficulties. *Journal of Experimental Child Psychology*, 87, 85–106.
- Gathercole, S. E. (1999). Cognitive approaches to the development of short-term memory. *Trends in Cognitive Science*, 6, 76–82.
- Gunn, P., & Crombie, M. (1996). Language and speech. In B. Stratford, & P. Gunn (Eds.), *New approaches to Down syndrome* (pp. 249–267). London: Cassell.
- Hartman, E., Houwen, S., Scherder, E., & Visscher, C. (2010). On the relationship between motor performance and executive functioning in children with intellectual disabilities. *Journal of Intellectual Disability Research*, 54, 468–477.
- Hasselhorn, M., & Mähler, C. (2007). Phonological working memory of children in two German special schools. *International Journal of Disability, Development and Education*, 54, 225–244.
- Henry, L. A., & MacLean, M. (2002). Working memory performance in children with and without intellectual disabilities. *American Journal on Mental Retardation*, 107, 421–432.
- Henry, L. A., & MacLean, M. (2003). Relationships between working memory, expressive vocabulary and arithmetical reasoning in children with and without intellectual disabilities. *Educational and Child Psychology*, 20, 51–64.
- Henry, A. L., & Winfield, J. (2010). Working memory and educational achievement in children with intellectual disabilities. *Journal of Intellectual Disability Research*, 54, 354–365.
- Henry, L. A. (2001). How does the severity of a learning disability affect working memory performance? *Memory*, 9, 233–247.
- Henry, L. A. (2010). The episodic buffer in children with intellectual disabilities: An exploratory study. *Research in Developmental Disabilities*, 31, 1609–1614.
- Hitch, G. J., & McCauley, E. (1991). Working memory in children with specific arithmetic learning difficulties. *British Journal of Psychology*, 82, 375–386.
- Hulme, C., & Mackenzie, S. (1992). Working memory and severe learning difficulties. Hillsdale, NJ: Erlbaum.

- Hulme, C., Maughan, S., & Brown, G. D. (1991). Memory for familiar and unfamiliar words: Evidence for a long-term memory contribution to short-term memory span. *Journal of Memory and Language*, 30, 685–701. [http://dx.doi.org/10.1016/0749-596X\(91\)90032-F](http://dx.doi.org/10.1016/0749-596X(91)90032-F)
- Jarrold, C., & Hall, D. (2013). The development of rehearsal in verbal short-term memory. *Child Development Perspectives*, 7, 182–186.
- Jarrold, C., Baddeley, A. D., & Hewes, A. K. (2000). Verbal short-term memory deficits in Down syndrome: A consequence of problems in rehearsal? *Child Journal of Psychology and Psychiatry*, 41, 223–244.
- Jarrold, C., Baddeley, A. D., & Phillips, C. (2007). Long-term memory for verbal and visual information in Down syndrome and Williams syndrome: Performance on the Doors and People test. *Cortex*, 43, 233–247.
- Kane, M. J., & Engle, R. W. (2002). The role of prefrontal cortex in working memory capacity, executive attention, and general fluid intelligence: An individual-differences perspective. *Psychonomic Bulletin and Review*, 9(4), 637–671. OenURL Query Data [Medline].
- Lanfranchi, S., Cornoldi, C., & Vianello, R. (2002). Working memory deficits in individuals with and without mental retardation. *Journal of Cognitive Education and Psychology*, 2, 301–312.
- Lanfranchi, S., Carretti, B., Spanò, G., & Cornoldi, C. (2009). A specific deficit in visuo-spatial simultaneous working memory in Down syndrome. *Journal of Intellectual Disability Research*, 53, 474–483.
- Lanfranchi, S., Cornoldi, C., Drigo, S., & Vianello, R. (2009). Working memory in individuals with fragile X syndrome. *Child Neuropsychology*, 15, 105–119.
- Laws, G. (2002). Working memory in children and adolescents with Down syndrome: Evidence from a colour memory experiment. *Journal of Child Psychology and Psychiatry*, 43, 353–364.
- Levén, A., Lyxell, B., Andersson, J., Danielsson, H., & Rönnerberg, J. (2008). Prospective memory, working memory, retrospective memory and self-rated memory performance in persons with intellectual disability. *Scandinavian Journal of Disability Research*, 10(3), 147–165. <http://dx.doi.org/10.1080/15017410802144444>
- Levén, A., Lyxell, B., Andersson, J., Danielsson, H., & Rönnerberg, J. (2011). The relationship between prospective memory, working memory and self-rated memory performance in individuals with intellectual disability. *Scandinavian Journal of Disability Research*, 13(3), 207–223. <http://dx.doi.org/10.1080/15017419.2011.596649>
- Levy, Y., & Antebi, V. (2004). Word reading and reading-related skills in Hebrew-speaking adolescents with Williams syndrome. *Neurocase*, 10, 444–451.
- Levy, Y. (2011). IQ predicts word decoding skills in populations with intellectual disabilities. *Research in Developmental Disabilities*, 32, 2267–2277.
- Lifshitz, H., & Katz, Y. J. (2009). Religious concepts among individuals with intellectual disability: A comparison between adolescents and adults. *European Journal of Special Needs Education*, 24, 183–201.
- Lifshitz, H., Tzuriel, D., Weiss, I., & Tzemach, M. (2010). New model of mapping difficulties in solving analogical problems among adolescents and adults with intellectual disability. *Research in Developmental Disabilities*, 32, 326–344.
- Lifshitz-Vahav, H., & Vakil, E. (2014). Taxonomy of moderators that governed explicit memory in individuals with intellectual disability: Integrative research review. *Journal of Applied Research in Memory and Cognition*, 3, 101–119. <http://dx.doi.org/10.1016/j.jarmac.2014.03.001>
- Lifshitz-Vahav, H. (2015). Compensation Age Theory (CAT): A monograph on the effect of chronological age in a population of intellectual disability. *Education and Training in Autism and Developmental Disabilities*, 50, 142–154.
- Logie, R. A., & Pearson, D. G. (1997). The inner eye and the inner scribe of visuo-spatial working memory: Evidence from developmental fractionation. *European Journal of Cognitive Psychology*, 9, 241–257.
- Logie, R. H. (1986). Visuo-spatial processing in working memory. *Quarterly Journal of Experimental Psychology A: Human Experimental Psychology*, 38, 229–247. <http://dx.doi.org/10.1080/14640748608401596>
- Maehtler, C., & Schuchardt, K. (2009). Working memory functioning in children with learning disabilities: Does intelligence make a difference? *Journal of Intellectual Disability Research*, 53, 3–10.
- McCarthy, D. (1972). *McCarthy scales of children's abilities*. New York: Psychological Corporation.
- Merighi, J., Edison, M., & Zigler, E. (1990). The role of motivational factors in the functioning of mentally retarded individuals. In R. M. Hodapp, J. A. Burack, & E. Zigler (Eds.), *Issues in the developmental approach to mental retardation* (pp. 114–134). Cambridge: Cambridge University Press.
- Miller, J. F. (1992). Development of speech and language in children with Down syndrome. In I. T. Lotty, & E. E. McCoy (Eds.), *Down syndrome: Advances in medical care* (pp. 39–50). New York: Wiley-Liss.
- Millner, B. (1971). Interhemispheric differences in the localization of psychological processes in man. *British Medical Bulletin*, 3, 272–277.
- Miyake, A., Friedman, N. P., Rettinger, D. A., Shah, P., & Hegarty, P. (2001). How are visuo-spatial working memory, executive functioning and spatial abilities related? A latent-variable analysis. *Journal of Experimental Psychology: General*, 130, 621–640.
- Nash, H., & Heath, J. (2011). The role of vocabulary: working memory and inference making ability in reading comprehension in Down syndrome. *Research in Developmental Disabilities*, 32, 1782–1789.
- Numminen, H., Service, E., & Ruoppila, I. (2002). Working memory: Intelligence and knowledge base in adult persons with intellectual disability. *Research in Developmental Disabilities*, 23, 105–177.
- Numminen, H., Lehto, J. E., & Ruoppila, I. (2001). Tower of Hanoi and working memory in adult persons with intellectual disability. *Research in Developmental Disabilities*, 22, 373–387.
- Numminen, H., Service, E., Ahonen, T., & Ruoppila, I. (2001). Working memory and everyday cognition in adults with Down's syndrome. *Journal of Intellectual Disability Research*, 45, 157–168.
- O'Hearn, K., Courtney, S., Street, W., & Landau, B. (2009). Working memory impairment in people with Williams syndrome: Effects of delay, task and stimuli. *Brain and Cognition*, 69, 495–503.
- Palladino, P., Cornoldi, C., De Beni, R., & Pazzaglia, F. (2001). Working memory and updating processes in reading comprehension. *Memory and Cognition*, 29, 344–354.
- Pazzaglia, F., & Cornoldi, C. (1999). The role of distinct components of visuo-spatial working memory in the processing of texts. *Memory*, 7, 19–41.
- Perez, L. A., Peynircioğlu, Z. F., & Blaxton, T. A. (1998). Developmental differences in implicit and explicit memory performance. *Journal Experimental Child Psychology*, 70, 167–185.
- Piaget, J. (1970). *Piaget's theory*. Tel Aviv: Hapoalim [in Hebrew].
- Pickering, S., & Gathercole, S. E. (2001). *Working memory test battery for children (WMTB-C)*. New York: The Psychological Corporation.
- Poloczek, S., Büttner, G., & Hasselhorn, M. (2014). Phonological short-term memory impairment and the word length effect in children with intellectual disabilities. *Research in Developmental Disabilities*, 35, 455–462.
- Porter, M. A., Coltheart, M., & Langdon, R. (2007). The neuropsychological basis of hypersociability in Williams and Down syndrome. *Neuropsychologia*, 45, 2839–2849.
- Randolph, J. J. (2009). A guide to writing dissertation literature review. *Practical Assessment, Research & Evaluation*, 14, 1–13.
- Raven, J. C., Court, J. H., & Raven, J. (1992). *Standard progressive matrices*. Oxford: Oxford Psychologist Press.
- Reed, T. (1996). Analogical reasoning in participants with autism, retardation, and normal development. *Journal of Development and Physical Disabilities*, 8, 61–76.
- Roediger, H. L. (2008). Relativity of remembering: Why laws of memory vanished? *Annual Review of Psychology*, 59, 225–254.
- Roid, C. H., & Miller, L. G. (1997). *Leiter International Performance Scale-Revised: Examiner's Manual*. Wood Dale, IL: Stoeling.
- Rondal, J. A. (1996). Oral language in Down's syndrome. In J. Perera, L. Nadel, & A. Comblain (Eds.), *Down's syndrome: psychological, psychobiological and socio-educational perspectives* (pp. 99–117). London: Whurr.
- Rosenquist, C., Conners, F. A., & Roskos-Ewoldsen, B. (2003). Phonological and visuo-spatial working memory in individuals with intellectual disability. *American Journal on Mental Retardation*, 108, 403–413.
- Schuchardt, K., Gebhardt, M., & Mähler, C. (2010). Working memory functions in children with different degrees of intellectual disability. *Journal of Intellectual Disability Research*, 54, 346–353.

- Schuchardt, K., Mähler, C., & Hasselhorn, M. (2011). Functional deficit in phonological WM in children with intellectual disabilities. *Research in Developmental Disabilities*, 32, 1934–1940.
- Service, E. (1998). The effect of word length on immediate serial recall depends on phonological complexity, not articulatory duration. *Quarterly Journal of Experimental Psychology A: Human Experimental Psychology*, 51, 283–304. <http://dx.doi.org/10.1080/713755759>
- Torraco, R. (2005). Writing integrative literature reviews: Guidelines and examples. *Human Resource Development Review*, 4, 356–367.
- Van der Molen, M. J., Van Luit, J. E. H., Jongmans, M. J., & Van der Molen, M. W. (2007). Verbal working memory in children with mild intellectual disabilities. *Journal of Intellectual Disability Research*, 51, 162–169.
- Van der Molen, M. J., Van Luit, J. E. H., Jongmans, M. J., & Van der Molen, M. W. (2009). Memory profiles in children with mild intellectual disabilities: Strengths and weaknesses. *Research in Developmental Disabilities*, 30, 1237–1247.
- Van der Molen, M. J., Van Luit, J. E. H., Van der Molen, M. W., & Jongmans, M. J. (2010). Every day memory and working memory in adolescents with mild intellectual disabilities. *American Journal on Intellectual and Developmental Disabilities*, 113, 207–217.
- Vianello, R., & Marin, M. L. (1997). *Operazioni logiche e conservazione. Dal pensiero intuitivo al pensiero operatorio concreto: prove per la valutazione del livello di sviluppo*. Bergamo: Junior.
- Vicari, S., Carlesimo, G. A., & Caltagirone, C. (1995). Short-term memory in persons with intellectual disabilities and Down syndrome. *Journal of Intellectual Disability Research*, 39, 532–537.
- Vicari, S., Bellucci, S., & Carlesimo, G. A. (2003). Visual and spatial working memory dissociation: Evidence from Williams syndrome. *Developmental Medicine and Child Neurology*, 45, 269–273.
- Vicari, S., Bellucci, S., & Carlesimo, G. A. (2005). Visual and spatial long term memory: Differential pattern of impairments in Williams and Down Syndromes. *Developmental Medicine and Child Neurology*, 47, 305–311.
- Visu-Petra, L., & Benga, O. (2007). Visual-spatial processing in children and adolescents with Down's syndrome: A computerized assessment of memory skills. *Journal of Intellectual Disability Research*, 51, 942–952.
- Vygotsky, L. (1986). *Thought and language*. Cambridge, MA: The Massachusetts Institute of Technology Press.
- Wang, P. P., & Bellugi, U. (1994). Evidence from two genetic syndromes for a dissociation between verbal and visual-spatial short-term memory. *Journal of Clinical and Experimental Neuropsychology*, 16, 317–322.
- Wechsler, D. (1967). *Manual for the Wechsler preschool and primary scale of intelligence*. New York: The Psychological Corporation.
- Wechsler, D. (1991). *Manual for the wechsler intelligence scale for children—WISC-R-III* (3rd ed.). New York: The Psychological Corporation.
- Willner, P., Bailey, R., Parry, R., & Dymond, S. (2010). Evaluation of the ability of people with intellectual disabilities to 'weigh up' information in two tests of financial reasoning. *Journal of Intellectual Disability Research*, 54, 380–391.
- Zigler, E., & Balla, D. (Eds.). (1982). *Mental retardation. The developmental-difference controversy*. Hillsdale, NJ: Lawrence Erlbaum.