



Journal of Clinical and Experimental Neuropsychology

ISSN: 1380-3395 (Print) 1744-411X (Online) Journal homepage: https://www.tandfonline.com/loi/ncen20

Which verbal fluency measure is most useful in demonstrating executive deficits after traumatic brain injury?

Gitit Kavé , Eyal Heled , Eli Vakil & Eugenia Agranov

To cite this article: Gitit Kavé, Eyal Heled, Eli Vakil & Eugenia Agranov (2011) Which verbal fluency measure is most useful in demonstrating executive deficits after traumatic brain injury?, Journal of Clinical and Experimental Neuropsychology, 33:3, 358-365, DOI: 10.1080/13803395.2010.518703

To link to this article: <u>https://doi.org/10.1080/13803395.2010.518703</u>



Published online: 05 Nov 2010.

|--|

Submit your article to this journal 🖸





View related articles



Citing articles: 7 View citing articles 🗹

Which verbal fluency measure is most useful in demonstrating executive deficits after traumatic brain injury?

Gitit Kavé¹, Eyal Heled^{2,3}, Eli Vakil⁴, and Eugenia Agranov^{3,5}

¹Department of Education and Psychology, The Open University, Ra'anana, Israel

²Hebrew University, Jerusalem, Israel

³Sheba Medical Center, Ramat Gan, Israel

⁴Department of Psychology and Leslie and Susan Gonda (Goldschmied) Multidisciplinary Brain

Research Center, Bar Ilan University, Ramat Gan, Israel

⁵Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

This paper examines switching and clustering in phonemic and semantic fluency tasks in individuals with traumatic brain injury (TBI). Fluency tasks were administered to 30 Hebrew-speaking patients with TBI and 30 age-matched control participants. Significant group differences were found in total output, number of switches, and number of clusters on both tasks, but not in mean cluster size. Unlike prediction, z scores of the number of semantic switches and clusters were lower than the equivalent z scores on the phonemic test. Results highlight the executive component of semantic fluency and the importance of using this task when assessing cognitive functioning after TBI.

Keywords: Verbal fluency; Traumatic brain injury; Switching and clustering; Executive functions; Language testing; Hebrew.

Tests of verbal fluency (also called controlled word association tests) are commonly used in neuropsychological assessment of individuals with brain damage (Henry, Crawford, & Phillips, 2004), contributing to the evaluation of executive functioning. These tests ask individuals to generate as many different words as possible that follow a certain rule in a limited time frame, measuring the ability to search lexical stores, to retrieve information from semantic memory, and to switch cognitive set (Henry & Crawford, 2004a, 2004b; Kavé, 2005; Kavé, Avraham, Kukulansky-Segal, & Herzberg, 2007; Kavé, Kigel, & Kochva, 2008; Troyer, 2000). It has been suggested that phonemic fluency (i.e., retrieving words that begin with a certain letter) might be more impaired in individuals with frontal brain damage (Rosser & Hodges, 1994: Trover, Moscovitch, Winocur, Alexander, & Stuss, 1998), whereas semantic fluency (i.e., retrieving words

that belong to a given semantic category) might be more impaired in individuals with temporal brain damage. Therefore, some authors believe that the phonemic task should be more sensitive to traumatic brain injury (TBI) than the semantic fluency task (Capitani, Rosci, Saetti, & Laiacona, 2009; Juardo, Mataro, Verger, Bartumeus, & Junque, 2000). However, not all studies support this conclusion (Henry & Crawford, 2004a, 2004b). The current research conducts a quantitative and qualitative analysis of fluency variables, aiming to elucidate the cognitive components most sensitive to TBI.

Previous research on individuals with brain injury has demonstrated that their word production on fluency tests is limited (Axelrod, Tomer, Fisher, & Aharon-Peretz, 2001; Bittner & Crowe, 2006, 2007; Busch, McBride, Curtiss, & Vanderploeg, 2005; Capitani et al., 2009; Henry & Crawford, 2004b; Juardo et al., 2000).

This study was partially supported by a grant from the Rehabilitation Center for Veterans after traumatic brain injury (TBI) in Jaffa, Israel, given to the third author. We wish to thank Ayelet Avraham and Reut Lerer for their help in coding the data. There are no financial or other relationships that could be interpreted as a conflict of interests.

Address correspondence to Gitit Kavé, Department of Education and Psychology, The Open University, 1 University Road, P.O. Box 808, Ra'anana 43107, Israel (E-mail: gkave@012.net.il).

A meta-analysis of studies of TBI patients shows that the magnitude of the deficit cannot be accounted for by a failure to match patients and control participants on premorbid IQ, current verbal IQ, education, or psychomotor speed (Henry & Crawford, 2004b). In addition, impairments in word finding are not the principal cause of the reduction in fluency output, as patients with and without naming difficulties demonstrate equally deficient fluency (Bittner & Crowe, 2006). Instead, it is assumed that patients with TBI are impaired on fluency tests due to executive dysfunction, most notably due to difficulties in self-generative behavior (Busch et al., 2005). According to Capitani et al. (2009) as well as Juardo et al. (2000). TBI leads to greater deficits in phonemic rather than semantic fluency, probably due to its impact on the frontal lobes. However, TBI most often leads to diffuse damage in frontal as well as temporal brain regions (Vakil, 2005), and hence both fluency tasks might be impaired in individuals with TBI.

Word retrieval on all verbal fluency tests depends on lexical knowledge as well as on effective search processes that require set shifting (Troyer, Moscovitch, & Winocur, 1997). A meta-analysis of studies of persons with Alzheimer's disease, who have severe lexicalconceptual disorders, demonstrated greater difficulties in semantic fluency than in phonemic fluency (Henry et al., 2004). Furthermore, focal temporal damage has been associated with a lesser deficit on phonemic fluency and a larger deficit on semantic fluency (Henry & Crawford, 2004a). Although persons with temporal lesions show a semantic fluency deficit, this deficit is often found in individuals with frontal lesions as well (Rogers, Ivanoiu, Patterson, & Hodges, 2006). It has been shown that persons with focal frontal lobe lesions produce significantly fewer words on phonemic fluency tests than do healthy controls, and also perform worse on that test than do persons with nonfrontal lesions (Alvarez & Emory, 2006). Yet, according to Henry and Crawford (2004a, 2004b), frontal brain damage, whether focal or diffuse, leads to a comparable impairment on both phonemic and semantic tests.

Some authors have used qualitative methods to examine the cognitive strategies underlying verbal fluency tasks (e.g., Fagundo et al., 2008; Kavé et al., 2007, 2008; Koren, Kofman, & Berger, 2005; Kosmidis, Vlahou, Panagiotaki, & Kiosseoglou, 2004; Lanting, Haugrud, & Crossley, 2009; Rosselli, Tappen, Williams, Salvatierra, & Zoller, 2009; Troyer et al., 1997; Woods et al., 2004). Instead of comparing the total number of words provided on the phonemic and semantic tasks, these analyses have focused on two components termed switching and clustering. According to Troyer et al. (1998), when generating words on the phonemic and semantic fluency tasks, participants produce clusters of phonemically or semantically related words, and once a subcategory is exhausted they switch to another subcategory. Thus, performance on these tasks relies on (a) an executive component (i.e., switching) responsible for strategic search, response initiation, monitoring, shifting, and flexibility; and (b) an associate component (i.e., clustering) that reflects the semantic organization of memory stores (Troyer, 2000; Troyer et al., 1997; Troyer et al., 1998).

It is further assumed that anterior brain regions play a more important role in switching than in clustering. Troyer et al. (1998) examined this hypothesis in persons with focal brain lesions, finding individuals with frontal lobe lesions to switch less frequently than healthy participants and to produce normal cluster size on both the phonemic and the semantic tasks. In contrast, individuals with temporal lobe lesions exhibited normal switching and clustering on the phonemic task, but were impaired in switching on the semantic task. Although persons with temporal lobe lesions showed no marked deficit in cluster size, those who had left temporal lesions produced smaller clusters than those who had right temporal lesions. This study suggested that phonemic clustering was less dependent on the integrity of lexical stores than was semantic clustering and that the most discriminating index among the patient groups was the number of switches on the phonemic fluency task, which was impaired only in persons with frontal lesions.

Additional evidence supporting the assumption that switching is an executive function, whereas clustering is more dependent on lexical abilities, especially within the semantic task, comes from research of various nonfocal neuropsychological disorders, such as dementia due to HIV (Woods et al., 2004), adults with attentiondeficit/hyperactivity disorder (ADHD; Tucha et al., 2005), multiple sclerosis (Troster et al., 1998), and depression (Fossati, Guillaume, Ergis, & Allilaire, 2003). On the other hand, individuals with Alzheimer's disease produce smaller clusters than normal (Epker, Lacritz, & Cullum, 1999; Fagundo et al., 2008; Troster et al., 1998). In addition, persons with schizophrenia, who suffer from disproportionate semantic fluency impairment relative to phonemic fluency (Bokat & Goldberg, 2003; Kremen, Seidman, Faraone, & Tsuang, 2003), show a disproportionate decrease in the number of clustered words (Bozikas, Kosmidis, & Karavatos, 2005).

Thus, the examination of switching and clustering can clarify the relative contribution of executive strategies and semantic stores to fluency performance in TBI. Since TBI primarily affects frontal regions, we predict that measures of executive functioning (e.g., the number of switches as well as the number of clusters) will be more impaired in this population than measures that represent semantic storage (e.g., cluster size). We want to examine whether executive deficits will be comparable on the phonemic and semantic tasks or more pronounced on one task than on the other. The aim of the current paper, then, is to determine whether qualitative measures of verbal fluency contribute to our understanding of the underlying mechanisms that lead to fluency deficits in TBI.

METHOD

Participants

Sixty native Hebrew speakers participated in the study, 30 in the TBI group and 30 in the control group, each

group consisting of 22 men and 8 women. The TBI group was recruited at the day treatment brain injury unit of the Sheba Medical Center, Tel Hashomer, Israel. Patients had moderate to severe TBI, with documented head injury as determined by radiological findings. Inclusion criteria consisted of 6-24 months post injury, a Glasgow Coma Scale (GCS) score of 3-12, loss of consciousness of at least 30 min, age between 18-35 years, and at least 12 years of education. Exclusion criteria included previous brain damage, psychiatric disorder prior to injury, combination of TBI and spinal cord injury, and aphasia. Table 1 summarizes background information for patient and control groups. Post-traumatic amnesia (PTA) was determined for every patient, operationally defined as the time when the patient was oriented to self, place, and time, remembered information presented by the examiner, and recognized three objects (as in Shores, Marosszeky, Sandanam, & Batchelor, 1986). All questions addressed at the patient had to be answered correctly for a three-day period, and PTA was counted up to the first of these three days. Verbal IO was available for 22 (73%) of the patients and was within normal limits in all cases.

The control group was selected from Kavé et al.'s (2007) sample to match the TBI group in age and gender. Healthy participants were recruited through places of employment, university classes, and word of mouth. Persons with a known history of learning disorders, psychiatric disturbances, neurological disease, or head trauma were not included in the study. No verbal IQ scores were available for the healthy participants. The two groups did not differ in age, t(58) = -0.164, *ns*, but the TBI group was slightly less educated than the control group, t(58) = -2.298, p < .05 (see Table 1), as expected by their age of injury. Participant recruitment in both groups was conducted in accordance with institutional research guidelines.

Procedure

Participants were asked to provide as many words as possible within 60 seconds on each of three letters (phonemic test) and three categories (semantic test). The phonemic fluency test was administered first and then the semantic fluency test, and the order of letters, as well as the order of semantic categories, was constant across participants. Responses were written verbatim, with errors or repetitions subsequently excluded from the total score. When a questionable response was provided, clarifications were invited at the end of the one-minute interval.

Phonemic fluency was assessed by obtaining the number of words generated in one minute for the letters bet (/b/), gimel (/g/), and shin $(/\check{s}/)$. Instructions were as follows: "I want you to say as many Hebrew words as possible that begin with a certain letter. You may say any word except for names of people and places, such as *Tomer* or *Tel Aviv*. Also, you should use different words rather than the same word with a different ending. For example, if you say *tapuz* ('orange'), don't also say *tapuzim* ('oranges'). If you say a verb, use the simplest form halax ('he went') and not halaxti ('I went') or holex ('he goes'). Please don't say words that are attached to other words, such as mi-shamayim ('from the sky') or la-kise ('to the chair')."

Semantic fluency was assessed by obtaining the number of words generated in one minute for each of the following three semantic categories: animals, fruits and vegetables, and vehicles. Fruits and vegetables were treated as one category in order to avoid the ambiguity between botanical definitions and common usage (as in "avocado"). It was specified that for the category of vehicles only types of transportation should be provided while brand names were unacceptable.

Scoring

When homophones were provided, the second mention was counted only if the participant pointed out the alternate meaning explicitly (i.e., ganal "camel," "repaid"). Words inflected in both masculine and feminine forms (e.g., gever-gveret "mister-mistress"; sus-susa "horsemare") were counted as one, whereas an animal and its offspring were counted as separate words (e.g., para "cow" and egel "calf"). Synonyms were counted as two (matos and aviron "airplane"). Names of subcategories on the semantic test (e.g., bird) were not given credit if specific items within that subcategory (e.g., dove, eagle)

 TABLE 1

 Sample characteristics by group

	TBI gr	oup (N = 30)	Control group $(N = 30)$			
	Range	Mean (SD)	Range	Mean (SD)		
Age (years)	19–35	25.47 (4.69)	19-35	25.67 (4.74)		
Education (years)	12-16	12.63 (1.33)	12-16	13.47 (1.48)		
Time post injury (months)	6–24	12.47 (5.39)				
GCS	3-12	6.00(2.44)				
LOC (days)	0.5-90	17.45 (19.01)				
PTA (days)	4-180	51.63 (44.55)				
VIQ (N = 22)	82-120	102 (11.62)				

Note. TBI = traumatic brain injury; GCS = Glasgow Coma Scale; LOC = loss of consciousness; PTA = post-traumatic amnesia; VIQ = Wechsler Adult Intelligence Scale–Third Edition (WAIS–III) Verbal IQ.

were also provided. Slang terms (e.g., *shluk* "sip"), as well as foreign words (e.g., bandana, gangster), were generally acceptable.

Clustering and switching

In line with Troyer et al.'s (1997) guidelines, repetitions and mistakes were included in the scoring of clustering and switching. An item that appeared in two clusters was coded in both. For example, the "cat" in "dog, cat, tiger, lion" was counted both as part of the cluster of pets and as part of the cluster of felines. In cases in which a small cluster was embedded within a larger cluster, only the larger cluster was counted. Thus, if farm birds were generated among other farm animals, only one cluster was counted (i.e., horse, cow, chicken, duck, turkey = 1 cluster). Semantic clusters generated within the phonemic task, as well as phonemic clusters generated within the semantic task, were not scored.

Phonemic clustering

A cluster was counted when two consecutive words shared the first consonant and vowel (*gezer-geshem*), shared the first and second consonant but differed in the vowel of the opening syllable (*gina-ganav*), rhymed (*shamayim-shinayim*), or included duplication (*barburbilbul*).

Semantic clustering

Where possible, subcategories were based on previous studies (Kosmidis et al., 2004; Troyer et al., 1997). Guidelines were formulated for consistency sake but flexibility was allowed for the coding of associated words that did not fall under the list of predefined clusters. On the Animal category, clusters were coded according to habitat, zoological family, and family relation, which were further classified into relevant subcategories. On the Fruit and Vegetable category, clusters were coded according to either fruits or vegetables, with subcategories further defined by season, botanical family, manner of eating, and so on. On the Vehicle category, clusters were coded according to land, water, or airborne means of transportations, with further classifications within land vehicles defined by common use (see Kavé et al., 2008, for further details of subcategories).

Four variables were derived for each fluency test on the basis of the aforementioned criteria:

Total fluency score

All words, excluding repetitions and errors, were summed across the three letters for the phonemic task and across the three categories for the semantic task.

Mean cluster size

Following Troyer et al. (1997), the number of words in a cluster was counted from the second word. That is, a cluster of two words was coded as 1, a cluster of three words was coded as 2, and so forth. A mean of all clusters of two words or more was computed for every person for each letter or semantic category. These means were then averaged across the three letters to yield the mean phonemic cluster size of each participant and across the three semantic categories to yield the mean semantic cluster size of each participant.

Number of switches

The number of switches between clusters of two words or more, between a cluster and a single word generated outside a cluster, and among those out-of-cluster single words (as in Troyer et al., 1997), was counted for every person for each letter and for each semantic category. Switches produced by each participant were summed across the three letters to yield the total phonemic number of switches score, and across the three semantic categories to yield the total semantic number of switches score.

Number of clusters

The number of clusters was counted separately, without single words, in order to examine participants' use of word association. As noted by Koren et al. (2005), the presence of single words may indicate that participants are in fact unable to utilize an associative strategy, and thus a measure that leaves out the single words is essential when focusing on the tendency to produce related words.

Inter-rater reliability

One of the raters who coded the data in Kavé et al. (2007) also coded all patient data. In order to calculate interrater reliability, another rater coded the responses generated by 10 participants with TBI. On the phonemic task, correlations between the two raters were r(8) = .91, p < .01, for mean cluster size, r(8) = .97, p < .01, for number of switches, and r(8) = .84, p < .01, for number of clusters. On the semantic task, correlations were r(8)= .97, p < .01, for mean cluster size, r(8) = .96, p < .01,for number of switches, and r(8) = .81, p < .01, for number of clusters. Control data were not coded again but rather taken from Kavé et al. (2007). Interrater reliability for control data was based on correlations between two judges who coded 30 protocols out of 100. All correlations between these two judges were high and significant at the .01 level (see Kavé et al., 2007, for exact details).

RESULTS

To rule out the possibility that the higher education level of the control group accounted for the current results, correlation analyses were conducted between years of education and fluency scores. Within the control group alone, only phonological cluster size was significantly related to education, r(28) = -.403, p < .05, with larger clusters in less educated individuals. The other seven correlations were not significant (p > .05). Within the TBI

Fluency test	Variable	TBI group (N=30)			Control group $(N = 30)$		
		Mean	SD	Range	Mean	SD	Range
Phonemic	Total 3 letters	25.80	8.21	9–40	45.50	10.87	27–68
	Mean cluster size	1.31	0.41	0.67 - 2.67	1.51	0.30	1.07-2.18
	Sum of switches	16.47	6.22	4-27	28.97	6.64	16-42
	Sum of clusters	6.47	2.65	1-13	12.40	4.83	4–26
Semantic	Total 3 categories	39.63	11.07	17-70	64.03	11.20	44-88
	Mean cluster size	1.58	0.32	0.97-2.22	1.80	0.34	1.28-2.45
	Sum of switches	20.87	5.91	8-33	33.60	4.80	26-41
	Sum of clusters	11.50	3.79	5-22	18.30	3.81	11-25

 TABLE 2

 Means, standard deviations, and range of scores of the eight fluency variables, by group

Note. Results are presented for all three letters together on the phonemic task and for all three categories together on the semantic task. TBI = traumatic brain injury.

group, none of the eight measures was significantly correlated to education. When correlations were examined for the two groups together, the number of switches on the phonological task was significantly correlated with education, r(58) = .378, p < .05, with more switches produced by more highly educated individuals. No other correlations were statistically significant, and therefore we decided not to enter years of education as a covariate in the analyses described below.

Table 2 displays means and standard deviations of fluency performance for each group. Because each variable was scored on a different scale, we chose not to run a factorial mixed design with repeated measures comparisons that would look at all variables together. Instead, raw scores were analyzed by eight independent-samples t tests, conducted for each variable separately. This analysis allows us to compare groups rather than focus on the comparison of performance across tasks, as production on the phonemic task is expected to be lower for each individual than production on the semantic task (Kavé, 2005). A Bonferroni correction was used to adjust the level of significance, and each comparison was examined against a significance level of .00625 (.05/8). The TBI group produced significantly fewer words than did the control group on both the phonemic task, t(58) =-7.917, p < .001, Cohen's d' = -2.04, and the semantic task, t(58) = -8.490, p < .001, Cohen's d' = -2.19. No significant group differences were found for mean phonemic cluster size, t(58) = -2.083, p = .042, Cohen's d' = -0.53, or for mean semantic cluster size, t(58) = -2.637, p = .011, Cohen's d' = -0.67. Individuals with TBI produced significantly fewer phonemic switches than did healthy participants, t(58) = -7.522, p < .001, Cohen's d' = -1.88, as well as significantly fewer semantic switches, t(58) = -9.161, p < .001, Cohen's d' = -2.37. The TBI group also produced significantly fewer clusters than did the control group on both the phonemic task, t(58) =-5.897, p < .001, Cohen's d' = -1.52, and the semantic task, t(58) = -6.929, p < .001, Cohen's d' = -1.79. Note that the strongest effect size was found for the comparison of the number of semantic switches.

To determine which fluency task is most sensitive to TBI, we calculated the sensitivity and specificity of both

tasks. We thus compared scores in both groups to ageappropriate normative scores that did not include the healthy participants of the current study (Kavé, 2005), looking at total scores only as there are no normative data for the qualitative variables. We chose a cutoff score of 1.5 standard deviations (SDs) below the mean normative score and counted how many individuals in each group scored outside that range. This analysis showed that 19 patients scored below the 1.5-SD cutoff score on the phonemic test, setting the sensitivity of the test at 63%, whereas the semantic test showed sensitivity of 80%(i.e., 24 patients scored below the cutoff). Specificity was high for both tests: 96% for the phonemic test (i.e., 29 out of 30 healthy individuals scored above the cutoff), and 93% for the semantic test (that is, 28 out of 30 healthy individuals scored above the cutoff).

Next we computed z scores for each participant on all fluency variables using the mean and standard deviation of the control group in the current study. This method of standardization was selected over using the mean and standard deviation of all participants together so that it would be possible to compare standardized scores across tasks (standardizing scores on the basis of data from all participants together would result in a mean of zero in all tasks). This analysis allows us to determine deviation from normal performance, as is done in clinical setting as well as in previous research (for instance, Vakil, Blachstein, Rochberg, & Vardi, 2004). Table 3 presents the means of z scores for each of the eight variables. Setting a cutoff score of 1.5 SD below the mean of the control group identified 26 patients (87%) as impaired on the total semantic score and 21 patients (70%) as impaired on the total phonemic score (see Table 3). Similar asymmetries between the semantic and phonemic tasks were found for the number of switches (83% vs. 70% patients below cutoff, respectively), as well as the number of clusters (60% vs. 40% patients below cutoff, respectively). Mean z scores of cluster size in the TBI group indicated no impairment on either fluency test.

After standardizing all scores within the TBI group according to the mean score of the control group in the current study, we plotted z scores by variable in order to compare impairments across fluency tasks using the same

	Variable	TPL - secre				Control 7 soore		
Test			I BI Z SCORE				Control 2 score	
		Mean	SD	Range	z < -1.5	Range	<i>z</i> < –1.5	
Phonemic	Total 3 letters	-1.84	0.77	-3.41 to -0.51	N = 21	-1.73 to 2.10	N = 1	
	Mean cluster size	-0.67	1.41	-2.90 to 4.00	N = 9	-1.53 to 2.30	N = 1	
	Sum of switches	-1.91	0.95	-3.82 to -0.30	N = 21	-1.99 to 2.00	N = 4	
	Sum of clusters	-1.25	0.56	-2.40 to 0.13	N = 12	-1.77 to 2.86	N = 2	
Semantic	Total 3 categories	-2.22	1.01	-4.27 to 0.54	N = 26	-1.82 to 2.18	N = 2	
	Mean cluster size	-0.67	0.97	-2.52 to 1.27	N = 6	-1.59 to 1.97	N = 2	
	Sum of switches	-2.70	1.25	-5.42 to -0.13	N = 25	-1.61 to 1.57	N = 1	
	Sum of clusters	-1.82	1.01	-3.56 to 0.99	N = 18	-1.95 to 1.79	N = 2	

 TABLE 3

 The z scores of the TBI group relative to the mean scores of the control group

Note. The *z* scores were calculated on the basis of the mean and standard deviation of the raw data in the control group. Thus, the mean of *z* scores within the control group was 0 for all variables, and the standard deviation of *z* scores was always 1 in that group. TBI = traumatic brain injury.



Figure 1. Means of z scores in TBI group relative to those of control group in each fluency variable, by test.

scale. Figure 1 shows that the mean standardized score of the number of semantic switches was the lowest of all z scores. In general, standardized scores on the semantic test were lower for all measures than standardized scores on the phonemic test, except for mean cluster size, which was equal across both tasks.

Finally, to determine whether the phonemic or the semantic test best discriminated between groups, we conducted four repeated measures analyses of variances on standardized scores, with task (phonemic, semantic) as the within-subject variable and group (TBI, control) as the between-subject variable. As variables are inherently dependent on each other, so that by definition the number of switches or clusters is defined by the number of words, we chose to conduct a separate analysis for each variable. An analysis that compared standardized total scores found a significant main effect of group, F(1, 58) = $81.949, p < .001, \eta^2_p = .586$, but no significant main effect of task, F(1, 58) = 3.200, ns, and no significant interaction F(1, 58) = 3.212, ns. An analysis that compared standardized scores of number of switches found a significant main effect of group, F(1, 58) = 99.938, p < .001, η^2_{p} = .633, a significant main effect of task, F(1, 58) = 6.742, p < .05, $\eta^2_p = .104$, and a significant interaction F(1, 58)= 6.812, p < .05, $\eta^2_{p} = .105$. As can be seen in Table 3, the difference between the mean z score of phonemic switches and the mean z score of semantic switches was greater in the TBI group than it was in the control group,

showing a greater deficit in switching on the semantic task. An analysis that compared standardized scores of the number of clusters found a significant main effect of group, F(1, 58) = 61.121, p < .001, $\eta^2_p = .513$, a significant main effect of task, $F(1, 58) = 4.369, p < .05, \eta^2_{p}$ = .070, and a significant interaction, F(1, 58) = 4.349, p < .05, $\eta^2_{p} = .070$. These results corroborate the results of the analysis of the number of switches, showing a greater deficit on the semantic than on the phonemic task within the TBI group. An analysis that compared standardized scores of cluster size found a significant main effect of group, F(1, 58) = 8.218, p < .05, $\eta^2_p = .124$, but no significant main effect of task, F(1, 58) = .004, ns, as well as no significant interaction, F(1, 58) = .002, ns. That is, patients made significantly smaller clusters than did control participants across tasks, but there was no difference between the size of clusters on the phonemic and the semantic tests.

DISCUSSION

Contrary to prediction, individuals with TBI demonstrated a greater deficit on the semantic fluency task than on the phonemic fluency task, yet their deficit reflects difficulties in switching rather than in clustering, thus attesting to an underlying executive dysfunction. Previous studies suggested that phonemic fluency might be more impaired in individuals with TBI than semantic fluency (Capitani et al., 2009; Juardo et al., 2000; Rosser & Hodges, 1994; Troyer et al., 1998), because the phonemic task presumably relies more heavily on frontal brain structures, whereas the semantic task relies more heavily on temporal brain regions. Nevertheless, this suggestion has not received consistent support in the literature, perhaps because TBI is not necessarily confined to frontal regions (Vakil, 2005). In fact, in their meta-analyses of earlier studies, Henry and Crawford (2004a, 2004b) showed that frontal brain damage leads to comparable impairment on both phonemic and semantic tests. Furthermore, Kavé et al. (2007) found that scores on both fluency tasks were equally associated with performance on a test of mental flexibility (the Homophone Meaning Generation Test, HMGT) through a common executive component.

Both fluency tasks involve strategic search, response initiation, monitoring, shifting, and flexibility, which can be measured through the number of switches (Troyer et al., 1997), as well as the number of clusters (Kavé et al., 2008; Koren et al., 2005), whereas the size of the clusters generated by participants reflects reliance on lexical stores. Indeed, patients with TBI are known to be impaired on a variety of executive tasks that include selfgenerative behavior, flexibility and set shifting, mental control, and self monitoring (Busch et al., 2005). It is not surprising, then, that our patients were found to be impaired on both fluency tasks, as well as on both variables that measured set shifting (i.e., number of switches and number of clusters), but no deficit in cluster size was seen in this group. Note, though, that the current results did not reveal entirely comparable deficit on both fluency tests. Instead we found that patients were more impaired on the semantic test than on the phonemic test.

Our findings suggest that the semantic test is more sensitive to TBI than is the phonemic test, when scores are compared either to normative data or to data of matched control participants. Moreover, the number of switches and the number of clusters were more affected by brain injury on the semantic task than on the phonemic task. In fact, the mean standardized score of the number of clusters on the phonemic task was within normal range. Importantly, the variable that led to the lowest standardized scores was the number of switches on the semantic task. It appears, then, that the semantic test is sensitive to the difficulties caused by TBI specifically because it taps into executive processes.

Note that phonemic fluency is most typically assessed through a sum score of responses to three letters (FAS in English; b, g, sh in Hebrew), yet this is not always the case for semantic fluency, which has been commonly evaluated with the single category of "animals" (e.g., Curtis, Thompson, Greve, & Bianchini, 2008; Epker et al., 1999; Fagundo et al., 2008; Fossati et al., 2003, among many others). It is possible that the current results reflect the fact that we used three semantic categories rather than just one. Indeed, when the category of animals alone was examined in the current study, the effect size of the group difference on this category was not only smaller than the effect found for the three semantic categories together (Cohen's d' = -1.93 for animals alone vs. Cohen's d' = -1.932.19 for all three categories), but also slightly smaller than the effect found for the three letters together (Cohen's d' = -1.93 for animals alone vs. Cohen's d' = -2.04 for all three letters). It has been suggested that test-retest reliability of the three-letter version is better than reliability for any one letter (Harrison, Buxton, Husain, & Wise, 2000), most likely because sampling a greater range of behavior improves the validity of results. It is safe to assume that sampling word output of three categories improves the validity of the semantic test relative to sampling one category alone, especially since the interindividual variance is large within both healthy and

brain-injured groups of participants. Thus, it could be the case that previous findings of a greater deficit in phonemic than in semantic fluency reflected in part the fact that the former was assessed over three letters (and three minutes), whereas the latter was assessed with only one category.

Several limitations of our study design must be acknowledged. Had we used other cognitive tests we could have better characterized the underlying cognitive processes leading to the observed impairment. Unfortunately, we administered no other executive tests to all participants and had no independent measures of processing speed. While these data were available for some patients, the information was available neither for all patients nor for the healthy participants. Thus, we cannot rule out the possibility that the differences recorded here reflect compromised speed of processing rather than a deficit in executive abilities. Note, however, that a decrease in processing speed would most probably lead to a comparable impact on both fluency tasks. Another limitation involves the selection of control participants. As the TBI patients underwent significant trauma, it would have been helpful to include other patients as controls (e.g., orthopedic patients) and thus to rule out the possibility that trauma by itself, rather than the head injury, was responsible for the reduced fluency performance.

Notwithstanding these limitations, our results can make a significant contribution to clinicians who work with the TBI population, by highlighting the importance of assessing semantic fluency in these patients and of doing so with more than one semantic category. To conclude, we show that semantic fluency is deficient in persons with TBI due to decreased switching, most likely reflecting impairment in flexibility that results from damage to the frontal as well as the temporal lobes.

> Original manuscript received 7 February 2010 Revised manuscript accepted 30 July 2010 First published online 5 November 2010

REFERENCES

- Alvarez, J. A., & Emory, E. (2006). Executive function and the frontal lobes: A meta-analytic review. *Neuropsychology Review*, 16, 17–42.
- Axelrod, B. N., Tomer, R., Fisher, T., & Aharon-Peretz, J. (2001). Preliminary analyses of Hebrew verbal fluency measures. *Applied Neuropsychology*, 8, 248–250.
- Bittner, R. M., & Crowe, S. F. (2006). The relationship between naming difficulty and FAS performance following traumatic brain injury. *Brain Injury*, 20, 971–980.
- Bittner, R. M., & Crowe, S. F. (2007). The relationship between working memory, processing speed, verbal comprehension and FAS performance following traumatic brain injury. *Brain Injury*, 21, 709–719.
- Bokat, C. E., & Goldberg, T. E. (2003). Letter and category fluency in schizophrenic patients: A meta-analysis. *Schizophrenia Research*, 64, 73–78.
- Bozikas, V. P., Kosmidis, M. H., & Karavatos, A. (2005). Disproportionate impairment in semantic verbal fluency in schizophrenia: Differential deficit in clustering. *Schizophrenia Research*, 74, 51–59.
- Busch, R. M., McBride, A., Curtiss, G., & Vanderploeg, R. D. (2005). The components of executive functioning in

traumatic brain injury. *Journal of Clinical and Experimental Neuropsychology*, 27, 1022–1032.

- Capitani, E., Rosci, C., Saetti, M. C., & Laiacona, M. (2009). Mirror asymmetry of category and letter fluency in traumatic brain injury and Alzheimer's patients. *Neuropsychologia*, 47, 423–429.
- Curtis, K. L., Thompson, L. K., Greve, K. W., & Bianchini, K. J. (2008). Verbal fluency indicators of malingering in traumatic brain injury: Classification accuracy in known groups. *The Clinical Neuropsychologist*, 22, 930–945. Epker, M. O., Lacritz, L. H., & Cullum, C. M. (1999).
- Epker, M. O., Lacritz, L. H., & Cullum, C. M. (1999). Comparative analysis of qualitative verbal fluency performance in normal elderly and demented populations. *Journal* of Clinical and Experimental Neuropsychology, 21, 425–434.
- Fagundo, A. B., Lópéz, S., Romero, M., Guarch, J., Marcos, T., & Salamero, M. (2008). Clustering and switching in semantic fluency: Predictors of the development of Alzheimer's disease. *Journal of the International Neuropsychological Society*, 23, 1007–1013.
- Fossati, P., Guillaume, L. B., Ergis, A. M., & Allilaire, J. F. (2003). Qualitative analysis of verbal fluency in depression. *Psychiatry Research*, 117, 17–24.
- Harrison, J. É., Buxton, P., Husain, M., & Wise, R. (2000). Short test of semantic and phonological fluency: Normal performance, validity and test–retest reliability. *British Journal of Clinical Psychology*, 39, 181–191.
- Henry, J. D., & Crawford, J. R. (2004a). A meta-analytic review of verbal fluency performance following focal cortical lesions. *Neuropsychology*, 18, 284–295.
 Henry, J. D., & Crawford, J. R. (2004b). A meta-analytic review
- Henry, J. D., & Crawford, J. R. (2004b). A meta-analytic review of verbal fluency performance in patients with traumatic brain injury. *Neuropsychology*, 18, 621–628.
- Henry, J. D., Crawford, J. R., & Phillips, L. H. (2004). Verbal fluency performance in dementia of the Alzheimer's type: A meta-analysis. *Neuropsychologia*, 42, 1212–1222.
- Juardo, M. A., Mataro, M., Verger, K., Bartumeus, F., & Junque, C. (2000). Phonemic and semantic fluencies in traumatic brain injury patients with focal frontal lesions. *Brain Injury*, 14, 789–795.
- Kavé, G. (2005). Phonemic fluency, semantic fluency, and difference scores: Normative data for adult Hebrew speakers. *Journal of Clinical and Experimental Neuropsychology*, 27, 690–699.
- Kavé, G., Avraham, A., Kukulansky-Segal, D., & Herzberg, O. (2007). How does the homophone meaning generation test associate with the phonemic and semantic fluency tests? A quantitative and qualitative analysis. *Journal of the International Neuropsychological Society*, 13, 424–432.
- Kavé, G., Kigel, S., & Kochva, R. (2008). Switching and clustering in verbal fluency tasks throughout childhood. *Journal of Clinical and Experimental Neuropsychology*, 30, 349–359.
- Koren, R., Kofman, O., & Berger, A. (2005). Analysis of word clustering in verbal fluency of school-aged children. *Archives* of Clinical Neuropsychology, 20, 1087–1104.
- Kosmidis, M. H., Vlahou, C. H., Panagiotaki, P., & Kiosseoglou, G. (2004). The verbal fluency task in the Greek population: Normative data, and clustering and switching strategies. *Journal of the International Neuropsychological Society*, 10, 164–172.

- Kremen, W. S., Seidman, L. J., Faraone, S. V., & Tsuang, M. T. (2003). Is there disproportionate impairment in semantic or phonemic fluency in schizophrenia? *Journal of the International Neuropsychological Society*, 9, 79–88.
- Lanting, S., Haugrad, N., & Crossley, M. (2009). The effect of age and sex on clustering and switching during speeded verbal fluency tasks. *Journal of the International Neuropsychological Society*, 15, 196–204.
- Rogers, T. T., Ivanoiu, A., Patterson, K., & Hodges, J. R. (2006). Semantic memory in Alzheimer's disease and the frontotemporal dementias: A longitudinal study of 236 patients. *Neuropsychology*, 20, 319–335.
- Rosselli, M., Tappen, R., Williams, C., Salvatierra, J., & Zoller, Y. (2009). Level of education and category fluency task among Spanish speaking elders: Number of words, clustering, and switching strategies. *Aging, Neuropsychology and Cognition*, 16, 721–744.
- Rosser, A., & Hodges, J. R. (1994). Initial letter and semantic category fluency in Alzheimer's disease, Huntington's disease, and progressive supranuclear palsy. *Journal of Neurology, Neurosurgery and Psychiatry*, 57, 1389–1394.
- Shores, E. A., Marosszeky, J. E., Sandanam, J., & Batchelor, J. (1986). Preliminary validation of a scale for measuring the duration of post-traumatic amnesia. *Medical Journal of Australia*, 144, 569–572.
- Troster, A. I., Fields, J. A., Testa, J. A., Paul, R. H., Blanco, C. R., Hames, K. A., et al. (1998). Cortical and subcortical influences on clustering and switching in the performance of verbal fluency tasks. *Neuropsychologia*, *36*, 295–304.
- Troyer, A. K. (2000). Normative data for clustering and switching on verbal fluency tasks. *Journal of Clinical and Experimental Neuropsychology*, 22, 370–378.
- Troyer, A. K., Moscovitch, M., & Winocur, G. (1997). Clustering and switching as two components of verbal fluency: Evidence from younger and older healthy adults. *Neuropsychology*, 11, 138–146.
- Troyer, A. K., Moscovitch, M., Winocur, G., Alexander, M. P., & Stuss, D. T. (1998). Clustering and switching on verbal fluency: The effects of focal-frontal and temporal-lobe lesions. *Neuropsychologia*, 36, 499–504.
- Tucha, O., Mecklinger, L., Laufkotter, R., Kaunzinger, I., Paul, G. M., Klein, H. E., et al. (2005). Clustering and switching on verbal and figural fluency functions in adults with attention deficit hyperactivity disorder. *Cognitive Neuropsychiatry*, 10, 231–248.
- Vakil, E. (2005). The effect of moderate to severe traumatic brain injury (TBI) on different aspects of memory: A selective review. *Journal of Clinical and Experimental Neuropsychology*, 27, 977–1021.
- Vakil, E., Blachstein, H., Rochberg, J., & Vardi, M. (2004). Characterization of memory impairment following closedhead injury in children using the Rey auditory verbal learning test (AVLT). *Child Neuropsychology*, 10, 57–66.
- Woods, S. P., Conover, E., Rippeth, J. D., Carey, C. L., Gonzalez, R., Marcotte, T. D., et al. (2004). Qualitative aspects of verbal fluency in HIV-associated dementia: A deficit in rule-guided lexical semantic search processes? *Neuropsychologia*, 42, 801–809.