

Neuropsychological Rehabilitation



An International Journal

ISSN: 0960-2011 (Print) 1464-0694 (Online) Journal homepage: www.tandfonline.com/journals/pnrh20

Memory outcomes following cognitive interventions in children with neurological deficits: A review with a focus on under-studied populations

Yael Schaffer & Ronny Geva

To cite this article: Yael Schaffer & Ronny Geva (2016) Memory outcomes following cognitive interventions in children with neurological deficits: A review with a focus on under-studied populations, Neuropsychological Rehabilitation, 26:2, 286-317, DOI: 10.1080/09602011.2015.1016537

To link to this article: https://doi.org/10.1080/09602011.2015.1016537



Published online: 02 Mar 2015.

ß	9

Submit your article to this journal 🕝



View related articles



View Crossmark data 🗹



Citing articles: 5 View citing articles 🖸

Memory outcomes following cognitive interventions in children with neurological deficits: A review with a focus on under-studied populations

Yael Schaffer and Ronny Geva

Department of Psychology, The Gonda Multidisciplinary Brain Research Center, Bar-Ilan University, Ramat Gan, Israel

(Received 19 October 2013; accepted 31 January 2015)

Given the primary role of memory in children's learning and well-being, the aim of this review was to examine the outcomes of memory remediation interventions in children with neurological deficits as a function of the affected memory system and intervention method. Fifty-seven studies that evaluated the outcome of memory interventions in children were identified. Thirty-four studies met the inclusion criteria, and were included in a systematic review. Diverse rehabilitation methods for improving explicit and implicit memory in children were reviewed. The analysis indicates that teaching restoration strategies may improve, and result in the generalisation of, semantic memory and working memory performance in children older than 7 years with mild to moderate memory deficits. Factors such as longer protocols, emotional support, and personal feedback contribute to intervention efficacy. In addition, the use of compensation aids seems to be highly effective in prospective memory tasks. Finally, the review unveiled a lack of studies with young children and the absence of group interventions. These findings point to the importance of future evidence-based intervention protocols in these areas.

Keywords: Paediatric; Rehabilitation; Memory; Restoration; Compensation.

Correspondence should be addressed to Prof. Ronny Geva, Department of Psychology, Head of the Developmental Neuropsychology Lab, The Gonda Brain Research Center, Bar-Ilan University, Ramat Gan, 5290002, Israel. E-mail: ronny.geva@biu.ac.il

The authors gratefully acknowledge Ms. Jessica Schreiber for her editorial input. This research was funded by an Israel Science Foundation [Grant #1518] awarded to Prof. Ronny Geva.

INTRODUCTION

Memory, the acquisition, retention and retrieval of information (Squire, 1992), is vital for daily functioning (Squire & Schacter, 2002), yet it is commonly affected by several different childhood central nervous system (CNS) disorders. These include genetically-based malformations, such as trisomy 21 (Conners, 2003; Jarrold, Baddeley, & Phillips, 1999; Jarrold, Purser, & Brock, 2012); acquired brain injury (ABI), such as traumatic brain injury (TBI) (Donders, 2007; Ewing-Cobbs, Fletcher, Levin, Iovino, & Miner, 1998; Souza, Braga, Filho, & Dellatolas, 2007), stroke (Kolk, Ennok, Laugesaar, Kaldoja, & Talvik, 2011) and brain tumours (Mulhern & Butler, 2004; Reeves et al., 2006); as well as various neurodevelopmental disorders, such as fetal alcohol syndrome (Mattson & Riley, 2006; Rasmussen, 2006), and human immunodeficiency virus (HIV) (Smith, Adnams, & Eley, 2008).

Memory is a multi-system construct (Sohlberg & Mateer, 2001). These childhood disorders affect memory functions in different ways as a function of the specific memory system that was affected (Manji, Pei, Loomes, & Rasmussen, 2009; Serra-Grabulosa et al., 2005; Ward, Shum, Wallace, & Boon, 2002). This review is based on a theoretical framework that differentiates between two major components of remembering: explicit and implicit memory systems, which typically refer to remembering past experiences (Schacter & Tulving, 1994); and a prospective memory system, which enables remembering to act in the future (Baddeley, 1997).

Explicit memory

Explicit or declarative memory refers to memories that can be consciously recalled (Baddeley & Hitch, 1994; Tulving, 1972; Willingham, Nissen, & Bullemer, 1989). Schacter and Tulving (1994) differentiate between three types of explicit memory: working memory (WM) (Baddeley, 2000; Baddeley & Hitch, 1994), semantic memory (SM) (Schacter & Tulving, 1994; Wilson & Glisky, 2009) and episodic memory (EM) (Tulving, 1972; Wilson & Glisky, 2009). Childhood CNS disorders often result in impairments in all three (Bangirana et al., 2009; Gorman, Barnes, Swank, Prasad, & Ewing-Cobbs, 2012; Jarrold et al., 2012; Kodituwakku, 2009; Mulhern & Butler, 2004, 2006; Pauly-Takacs, 2012; Serra-Grabulosa et al., 2005; Vargha-Khadem et al., 1997).

Implicit memory

The explicit memory system is complemented by the implicit memory system. Implicit memory is a non-declarative memory system that allows one to learn without conscious awareness, and it is composed of four memory types: priming, simple classical conditioning, non-associative learning and procedural memory (Schacter, 1987; Sohlberg & Mateer, 2001; Tulving, 1972; Tulving & Schacter, 1990). During the last decade many studies have demonstrated preserved implicit memory after acquired brain injury (ABI) in adults and paediatric populations (Vakil, 2005; Ward et al., 2002; Yeates & Enrile, 2005).

Prospective memory

The last memory component relevant for this review is prospective memory (PM). PM is the realisation of a delayed intention, and is commonly defined as remembering to perform an intended action in the future (Baddeley, 1997; Herrmann, Brubaker, Yoder, Sheets, & Tio, 1999; McDaniel & Einstein, 2007). Several CNS disorders, such as TBI, negatively affect PM in children (McCauley & Levin, 2004; Ward, Shum, McKinlay, Baker, & Wallace, 2007), thus rehabilitation for this type of memory seems warranted (McCauley & Pedroza, 2010).

This review explores studies that focus on implicit, explicit and PM improvement in children. As previously demonstrated, research shows consistently that memory deficits in one or more of these systems is a frequent consequence of childhood CNS disorders, and indicates that memory deficits have the potential to impede upon a child's daily functioning in his or her school (Hawley, 2004; Hawley, Ward, Magnay, & Mychalkiw, 2004), community (Van Heugten et al., 2006) and home environment (Kinsella et al., 1995; McCauley & Levin, 2004). In the last decade, cognitive rehabilitation has been frequently recommended for many paediatric patients (Ylvisker et al., 2005), although until recently, research on the effectiveness of memory rehabilitation techniques on paediatric patients with known neurological involvement was scarce (Laatsch et al., 2007; Limond & Leeke, 2005; Michaud, Rivara, Grady, & Reay, 1992). In the past eight years, however, four reviews of cognitive interventions in children (Laatsch et al., 2007; Limond & Leeke, 2005; Ross, Dorris, & McMillan, 2011; Tatla, Sauve, Jarus, Virji-Babul, & Holsti, 2014), one meta-analysis (Karch, Albers, Renner, Lichtenauer, & von Kries, 2013) and one review on the applicability of cognitive rehabilitation for children with ABI (Slomine & Locascio, 2009) have been conducted. These studies concentrated on paediatric populations with brain injury and ABI, and included relatively few studies specifically addressing memory deficits as the target for cognitive rehabilitation. In addition to these reviews, one meta-analysis on WM amenability to cognitive rehabilitation was recently conducted. This review mainly concerns children with attention deficit hyperactivity disorder (ADHD) (Melby-Lervåg & Hulme, 2013), but memory improvements in paediatric cohorts with known neuropathological aetiologies who are not primarily ADHD or autistic are markedly under-studied (Melby-Lervåg & Hulme, 2013). Based on the theoretical body of literature concerning memory systems, a systems-based analysis of intervention efficacy for the treatment of discrete memory systems in children with under-studied neurological disorders seems warranted.

Cognitive rehabilitation

Cognitive rehabilitation has been broadly defined as a systematic intervention designed to improve cognitive and/or behavioural difficulties following a pathological process in order to improve daily functioning (Axelrod et al., 2002). This intervention technique typically involves two essential modes: restoration (Sohlberg & Mateer, 2001) and compensation (Cicerone et al., 2000, 2005). Another approach that may support generalisation is behaviour management (van't Hooft & Norberg, 2009).

In the current context, it is important to note that the distinction between compensatory means as differentiated from restorative ones may have a heuristic value, but there may be some overlap and shared effects between the categories. Improved cognitive function, for example, may allow better compensation, and effective compensation may allow for some restoration of the underlying cognitive function. Nevertheless, this classification may potentially advance our understanding of the efficacy of memory rehabilitation (Sohlberg & Mateer, 2001), and therefore we use this classification here for heuristic purposes in order to delineate the efficacy of each intervention type for improving specific memory deficits.

The first mode of rehabilitation, restoration, aims to restore functions that have been lost. Restoration techniques suggest practising internal compensatory strategies in order to reduce a primary memory deficit (Sohlberg & Mateer, 2001). This is based on the idea that memory can be strengthened using memory drills analogous to a "mental muscle". Such strategies include rehearsal techniques (Belmont & Butterfield, 1969), chunking and mnemonics methods (Wilson & Moffat, 1992), and/or computerised brain training activities that aim to improve focused attention and the ability to ignore distractions (Berry, Zanto, Rutman, Clapp, & Gazzaley, 2009). Restoration can be accomplished explicitly by teaching specific strategies or implicitly by employing training.

The second mode of intervention may be a compensatory one that is based on the idea that planning abilities and external cues can be recruited to moderate the primary memory deficit. This is accomplished by offering effective means to circumvent the memory deficit, such as by encouraging the use of a diary, smartphone alerts, etc. (Cicerone et al., 2005, 2011). These external devices may help patients adjust to their memory deficit, and thus compensate for it in order to cope with everyday demands despite the cognitive impairment (Cicerone et al., 2005). A different compensatory approach aims to build upon preserved functions to support performance. An example of such an approach is errorless learning. This method was proposed by Wilson and colleagues, who found that since amnesic patients were unable to eliminate errors when learning new information and do not remember their learning experiences and the feedback that they receive due to a compromised explicit episodic memory system, they could learn more efficiently if, during learning, they were not required to produce potentially wrong responses (Wilson, Baddeley, Evans, & Shiel, 1994).

In addition to restoration and compensation, an important intervention method is behaviour management. Behaviour management uses operant techniques, such as conditioning or reinforcements (Domjan, 2003; McCauley, McDaniel, Pedroza, Chapman, & Levin, 2009), and social learning principles, such as parent management training (Braga, Da Paz, & Ylvisaker, 2005; Lindahl & Steneby, 2009), emotion-focused coping (Ramsden & Hubbard, 2002) and mediation, i.e., the interactions that parents have with children about their environment that promotes learning (Feuerstein, 1980), to achieve behaviour change. These methods have been thought of as a vital part of promoting learning directly and indirectly (van't Hooft & Norberg, 2009).

Across all intervention types, one of the major goals of memory intervention is generalisation, a process in which a learned strategy is used in response to situations other than the one for which it was practised (McKeough, Lupart, & Marini, 1995; Shadmehr & Moussavi, 2000). Nevertheless, while brain training programmes may improve performance on a specific subset of skills or tasks, the benefits may not generalise to other domains (Rabipour & Raz, 2012). Generalisation may be particularly challenging for patients with memory difficulties. Thus, it was examined in the current study as a function of intervention type.

The last aim of this review was to focus on young populations in order to gain more insight on brain plasticity and the role of age and neural maturation in paediatric rehabilitation. There is a debate focusing on the advantages of brain plasticity in younger ages (Bryck & Fisher, 2012; Kochanek, 2006; Wass, Scerif, & Johnson, 2012) versus the view that development and maturation may have both positive and negative effects on overall outcome (Giza & Prins, 2006). Therefore, the third aim of the analysis was to survey studies that focus on school aged children as well as younger children in order to gain greater insight on how different stages of maturation influence the ability to memorise a skill or a new set of information using different intervention methods (Bauer, DeBoer, & Lukowski, 2007; Carew & Magsamen, 2010; Jolles & Crone, 2012).

Hence, overall, this review's goals were to present available findings on memory rehabilitation outcomes in children with under-studied CNS disorders by classifying the different programmes into three inter-dependent categories: restoration, compensation or behaviour management; to delineate which memory aspects are the most frequently studied in that category; and to explore within each method category which memory system is responsive to the intervention, with age considered as a moderating factor.

METHODS

Literature search procedure

First, we surveyed the literature on paediatric memory rehabilitation by performing computer searches using PsycINFO (1967-2014), MEDLINE (1966-2014), American Psychological Association (APA) PsycNET (1967-2014), PsycBite 1966-2014, and Google Scholar databases. The searches used all possible combinations of the following terms and their synonyms (in English): rehabilitation/remediation/therapy, cognitive/neuro, paechild/children/adolescents, diatric/pediatric. memory/recollection/recall, acquired/traumatic brain injury/head injury, and training/treatment/management/intervention. The following search terms, covering all major causes of ABI and neurodevelopmental disorders, were also included: stroke/encephalitis/CNS, infection/oncology/fetal alcohol spectrum disorder/Down syndrome/spina bifida. All citations in all relevant articles were also identified and assessed. In an attempt to focus on the effect of cognitive rehabilitation in children suffering from memory deficits resulting from under-studied CNS disorders, studies focusing on ADHD and/or autism spectrum disorder (ASD) as primary inclusion criteria were not covered in the present review. The reason for this was that both ADHD and ASD are well studied, yet findings from these populations may dilute findings related to markedly under-studied populations who typically also present with attention and socio-communicative issues.

Figure 1 provides a flow diagram of the literature search strategy. The initial search yielded 1824 papers. Inclusion and exclusion criteria were then applied serially. Inclusion criteria were such that: (1) It was an intervention study that addressed memory intervention outcomes; and (2) the mean age of the participants was less than 20 years. Articles were excluded if: (1) The paper was not written in English; (2) the paper did not report original empirical data; (3) the study did not have at least one standardised outcome measure, and (4) the paper reported data on ADHD or ASD cohorts. Based on a review of study titles and these criteria, 1767 studies were excluded. The 57 remaining articles were then screened based on their abstracts using the stated inclusion and exclusion criteria, resulting in the exclusion of 10 more articles. A full article search was then performed on the remaining 47



Figure 1. A flow diagram of the papers' review process. ADHD = attention deficit hyperactivity disorder, ASD = autism spectrum disorder, CNS – central nervous system.

studies, and 13 more were excluded for reasons specified in the flow chart (Figure 1). At the end of this process, 34 articles were eligible for the review (detailed in Tables 1 and 2).

Methodological appraisal of included studies

Criteria for the assessment of articles were based on the guidelines of Consolidated Standards of Reporting Trials (CONSORT), with the added guidelines based on Ross and colleagues' (2011) paper (detailed in Appendix 1) resulting in 26 items. Each item was awarded a score of 1 if the criterion was met and 0 if it was not met or it was not possible to determine from the available information. Papers that met 75% of the methodological criteria specified were considered of "high" quality. Papers that were rated between 50 and 75% were deemed of "moderate" quality, and those achieving less than 50% were considered "low" quality. To assess the reliability of this tool, a second reviewer using the same system rated the final 34 papers. Overall percentage agreement was high (90%). Individual disagreement was resolved by discussion with the independent reviewer. This procedure helped address issues concerning the study's specific control conditions, how outcomes were assessed and treatment effect size (Tables 1 and 2).

Strategy type	Experimental Design	Quality rating*
One-to-one comprehensive	Randomised control study	High (76%)
One-to-one comprehensive	Randomised control trial	High (76%)
One-to-one comprehensive and environmental- familial support	Case study	Moderate (52%)
One-to-one comprehensive	Case study	Moderate (52%)
One-to-one comprehensive	Pre- and post- intervention assessment no control group	Moderate (47%)
	((Continued)

TABLE 1. Paediatric memory rehabilitation: Explicit memory methods

conditions Size (r)

Effect Follow

ир

**

Outcome

assessment

15 words list and

ROCFT, RALVT,

15 words list and

digit span test

digit span test

digit span

word list

ROCFT, digit span, One-to-one

Control

at the same age range

at the same

age range

to baseline

to baseline

to baseline performance

performance

performance

Memory

system

SM

SM

SM

SM

SM

Explicit Children

Explicit Children

Explicit None.

Explicit None.

Explicit None.

WM and with the ABI

WM and Comparison

WM and Comparison

WM and Comparison

WM and with the ABI

Intervention

AMAT-C

AMAT-C

AMAT-C

AMAT-C

AMAT-C

Type ofDur*

N insult freq

102

102

102

120

100

38 ABI

38 ABI

3 M

3 ABI

11-15 7 ABI

Authors

Van't

Hoolft

et al.

Van't

Hoolft

et al.

Van't

Hoolft

et al.

Van't

Hoolft

Sjo et al.

et al.

Year

2005

2007

2009

2003

2010

Age

9-16

9-17

9 - 14

10 - 16

MEMORY OUTCOMES FOLLOWING COGNITIVE INTERVENTIONS 293

TABLE 1
Continued

							Continu	cu					
Authors	Year	Age	Typ of N insu	e Dur* lt freq	Intervention	Memory system	Control conditions	Effect Size (r)	Follow up	Outcome assessment	Strategy type	Experimental Design	Quality rating*
Patel et al.	2009	7–19	12 C	15	Comprehensive programme including	Explicit WM and SM	None. Comparison to baseline performance	.15		CVLT-C	One-to-one comprehensive	Case series	Moderate (41%)
Butler and Copland	2002	6-18	31 C ar BT	d 50	Metacognitive Strategies and clustering skills CRP	Explicit WM and SM	Waiting list, or children who live far from cancer centre	.21		Digit span and sentence memory	Computerised	Control trial	Moderate (70%)
Butler et al	. 2008	6-17	161 CM	66	CRP	Explicit WM and SM	Waiting list	.15	**	Digit backward	Computerised	Randomised control trial	Moderate (70%)
										Stroop Color–Word Test Trail Making Test B			
Boivin et al.	2010	6-16	60 HIV	10	CCRT	Explicit WM and SM	Healthy controls	.27		Cogstate computerised neuropsychological test	Computerised	Randomised control trial	Moderate (73%)
Conners	2003	6-14	16 DS	108	Silent rehearsal	Explicit SM	Children with DS practising visual activities	.23		Digit span	One-to-one specific technique	Randomised cross over design	Moderate (64%)
Yerys et al	. 2003	11-15	6 SCE	6	Silent rehearsal and semantic organisation	Explicit SM	Children with SCD who received academic tutoring			Digit span, word list, semantic clustering	One-to-one specific technique	Randomised clinical trial	Moderate (47%)

King et al.	2007	8-16	11 SCD	104	Silent rehearsal and semantic clustering	Explicit SM	Children with SCD who received academic tutoring	.53		California Verbal Learning Test	One-to-one specific technique	Randomised control trial	Moderate (50%)
Oberg &	1998	18-19	2 TBI	10	Elaborative	Explicit	None.		**	% of learned word	One-to-one	Case studies -	Moderate
Turkustra				and 18	encoding	SM	to baseline performance			definitions	technique	intervention assessment	(47%)
Bangirana, et al. Giordan et al.	2009	10 ± 2.6	65 CM	16	CCRT	Explicit WM	Children with CM who received no training sessions	.28		One card learning and one back working memory task from Cogstate test	Computerised	Randomised trial	Moderate (70%)
Bangirana et al.	2011	5-12	61 CM	16	CCRT	Explicit WM	Children with CM who received no training sessions	29		Digit span	Computerised	Randomised trial	Moderate (70%)
Hardy et al.	2011	10-17	9 BT	24	CCRT	Explicit WM	None. Comparison to baseline performance	0.21	**	Digit span, letter- number sequencing	Computerised	Pilot with small sample	Moderate (54%)

(Continued)

Authors	Year	Age	N	Type of insult	Dur* freq	Intervention	Memory system	Control conditions	Effect Size (r)	Follow up	Outcome assessment	Strategy type	Experimental Design	Quality rating*
Bennet, Holmes, and Buckley	2013	7–12	21	DS	25	JCWMT	Explicit WM and SM	Waiting list	.11	**			Randomised cross over design	High (80%)
Loomes et al.	2008	4-11	33	FASD	4	Silent rehearsal	Explicit WM	Children with FASD who received no training sessions	33		Memory Test Battery for Children (WMTB-C)	One-to-one specific technique	Control trial	Moderate (62%)
Rankin and Hood	2005	9–15	2	М	3-9	Mnemonics	Explicit SM	None. Comparison to baseline performance			CMS, ROCFT	One-to-one specific technique	Case study	Low (29%)
Ponsford et al.	2001	6-15	211	TBI	1	Meta-memory Psycho- educational	Explicit WM	Children with minor injuries not involving the head	0		Assessment of memory and learning (WRAML) verbal scale score	Psycho- education	Randomised control trial	Moderate (68%)
Coyne et al.	in press	8-16	15	TBI	1	Retrieval practice	Explicit SM	Items practised with massed restudy, spaced restudy	.66		Number of correct responses across learning conditions	One-to-one	Randomised control trial	Moderate (69%)

TABLE 1 Continued

TBI = traumatic brain injury; BT = brain tumour; HIV = human immunodeficiency virus; CM = cerebral malaria; ABI = acquired brain injury; SCD = sickle cell disease; M = medulloblastoma; CM = childhood malignancy; FASD - fetal alcohol spectrum disorder; DS = Down syndrome; C = cancer. AMAT-C = Amsterdam Memory and Attention Training for children; CCRT = Captain's Log computerized cognitive rehabilitation therapy; CRP = Cognitive Remediation Program; JCWMT = Junior Cogmed Working Memory Training. WM = working memory; SM = semantic memory; LTM = long-term memory; NR = Not Reported. ROCFT = Rey Osterrieth Complex Figure Test; RALVT = Rey Auditory Learning Verbal Test; CVLT-C = California Verbal Learning Test-Children's version; WMTB-C = Working Memory Test Battery for Children; CMS = Child Memory Scale.

*Adapted from Ross & Dorris (2011), % of 26 quality criterions.

Authors	year	Age	N	Type of insult	Dur* freq	Intervention	Memory system	Control conditions	Effect Size (r)	Follow up	Outcome assessment	Strategy type	Experimental Design	quality rating*
Landis et al.	2006	6-18	34	TBI	14	Errorless learning	Immediate SM	Half of the items were taught the T&E method	15	**	Number of correct responses after the study trials	Errorless learning	Retrospective within-subjects concurrent treatment design	Moderate (65%)
Haslam, Bazen- Peters, and Wright	2012	11–16	30	ABI	1	Errorless learning	Immediate SM	15 healthy children	.79		Number of correct responses after the study trials	Errorless learning	Cross over design	Moderate (68%)
Van't Hoolft et al.	2005	9–16	38	ABI	102	AMAT-C	РМ	Children with the ABI at the same			RBMT	Compensation- comprehensive	Randomised control study	High (76%)
Van't Hoolft et al.	2007	9–17	38	ABI	102	AMAT-C	РМ	None. Comparison to baseline performance		**	RBMT	Compensation- comprehensive	Randomised control trial	High (76%)
Van't Hoolft et al.	2003	10-16	3	ABI	120	AMAT-C	РМ	None. Comparison to baseline performance			RBMT	Compensation- comprehensive	Case study	Moderate (52%)
Ho et al.	2011	11-17	15	ABI	6		PM	16 healthy control children	.25	**	ROCFT, CMQ and PMQ	Compensation	Case series	Moderate (52%)
Kerns and Thompson	1998	13	1	BT		Individualised memory notebook	PM	None		**	% accomplishing prospective memory tasks	Compensation	Case study – pre- and post- intervention assessment	Low (29%)
Wilson et al.	2009	8-17	12	TBI	7	Memory aids - Neuropage	• PM	Waiting list	.97	**	% accomplishing prospective memory tasks	Compensation	Randomised control trial	Moderate (56%)
Gillette and DePompe	2008	6-20	35	TBI	4	PDAs and smartphone:	PM	Waiting list versus two PDA groups versus planner			% accomplishing prospective memory tasks	Compensation	Control trial	Moderate (70%)

TABLE 2 Pediatric memory rehabilitation: Implicit memory methods

(Continued)

								Co	ntinued					
Authors	year	Age	N	Type of insult	Dur* freq	Intervention	Memory system	Control conditions	Effect Size (r)	Follow up	Outcome assessment	Strategy type	Experimental Design	quality rating*
McCauley	2009	6–19	84	TBI	2	Monetary incentive	РМ	Children with orthopaedic injury	.49		Amount of earned money	Behaviour management	Randomised cross over design	Moderate (64%)
McCauley	2010	6-19	119	TBI	2	Monetary incentive	РМ	Children with orthopaedic injury	.59–.44 for moderate TBI, –.12 for severe TBI		Amount of earned money	Behaviour management	Randomised cross over design	Moderate (64%)
McCauley et al	. 2011	7–16	115	TBI	2	Monetary incentive	РМ	Children with orthopaedic injury	.69 for moderate TBI, .22 for severe TBI	**	Amount of earned money	Behaviour management	Randomised cross over design	Moderate (64%)
Guillery- Girard et al.	2004	11-12	2		5	Errorless learning and vanishing cues	Immediate I SM				Number of correct responses after the study trials	Errorless learning	Case study	Moderate (60%)
Martins et al.	2006	7–9.5	2	BT and epilepsy	5	Errorless learning and vanishing cues	Immediate I SM	9 healthy controls			Number of correct responses after the study trials	Errorless learning versus vanishing cues	Case study	Moderate (60%)
Pauley- Takacs, Moulin, and Estlin	2012	15	1	BT	2	Errorless learning	Immediate SM	10 healthy controls	0.2	**	Number of correct responses after the study trials	Errorless learning	Case study	Moderate (65%)
Boivin et al.	2013	1.4-5	120	HIV	25	Parents' mediation	Explicit visual SM	60 healthy or HIV dyads who received health and nutrition curriculum	0.5		Color-Object Association Test (COAT) for memory	Behaviour management	Randomised control trial	High (84%)

TABLE 2

TBI = traumatic brain injury; BT = brain tumour; HIV = human immunodeficiency virus; ABI = acquired brain injury.

SM = semantic memory; PM = Prospective memory. AMAT-C = Amsterdam Memory and Attention Training for children; PDA = personal digital assistant; RBMT = Rivermead Behavioural Memory Test; ROCFT = Rey Osterrieth Complex Figure Test; CMQ = Child Memory Questionnaire; PMQ = Parent Memory Questionnaire; T & E =trial and error.

* Adapted from Ross & Dorris (2011), % of 26 quality criterions.

Recorded variables and coding

For the purposes of this review, each effect size was coded, along with several important pieces of information describing the population, the memory tasks, the intervention type and the study design.

Four studies employed several memory tasks and/or more than one measure of memory improvement. As a result, these reports included several effect sizes that were incorporated in the analysis wherever appropriate.

Individual effect sizes. We computed effect size using the formula described by Johnson and Eagly (2000), with the provided statistical information in each paper (e.g., means, standard deviations, student *t*-test, *F* test).

Characteristics of the studies

The literature search and inclusion screening yielded 34 articles, quality ratings from a systematic review standpoint (Ross et al., 2011) were conducted according to Standards of Reporting Trials (CONSORT) guidelines (see Appendix 1). Analysis indicated that four studies were considered high quality; 28 were rated as moderate quality and two studies were categorised as low quality. Importantly all included articles were highly commended from a clinical perspective.

Overall, this review included papers with data from 1375 youngsters with memory deficits who participated in intervention studies between the ages of 1.4 and 20 years (M age = 9.8 years, SD = 1.8 years). Most of the memory rehabilitation programmes (28 studies) were planned for and implemented with children older than 6 years. Only three studies (Bangirana et al., 2011; Boivin et al., 2013; Loomes, Rasmussen, Pei, Manji, & Andrew, 2008) included children younger than 6, pointing to a literature gap.

The childhood CNS disorders included in this review are: trisomy 21, acquired brain injury (ABI), traumatic brain injury (TBI), stroke, cerebral malaria, brain tumours, fetal alcohol syndrome and human immunodeficiency virus (HIV).

In most studies (n = 25), memory intervention programmes were implemented on a one-on-one basis (teacher, parent, or therapist), while in 10 studies memory intervention programmes were implemented with different computer programmes rather than with a social agent. Intervention in a group setting was not implemented in any of the studies (see Tables 1 and 2), pointing to a literature gap.

Twenty-two studies used various methods in order to improve explicit memory, and 16 investigated methods to improve implicit memory. Three studies used more than one intervention method in their protocol. We will

here describe intervention efficacy for each memory system, explicit, implicit and prospective memory systems, as a function of the primary intervention approach.

Explicit memory

Restorative techniques: Inner memory strategies. Restorative techniques for working memory (WM) usually involve chunking and mnemonics methods (Wilson & Glisky, 2009; Wilson & Moffat, 1992). Literature on cognitive rehabilitation of children's WM focuses mainly on two major approaches: a comprehensive strategic approach (one-on-one or computerised) and learning specific strategies (e.g., silent rehearsal or mnemonics). Both intend to promote inner compensation processes.

A comprehensive strategic approach was used in six studies, five of them examined the effect of the Amsterdam Memory and Attention Training for children AMAT-C on children's WM (Sjö, Spellerberg, Weidner, & Kihlgren, 2010; van't Hooft et al., 2005, 2007; van't Hooft, Andersson, Sejersen, Bartfai, & von Wendt, 2003; van't Hooft & Norberg, 2009) and the other examined a different rehabilitation protocol for children older than 7 years of age, teaching problem solving skills and compensatory strategies (Patel, Katz, Richardson, Rimmer, & Kilian, 2009).

All comprehensive protocols included diverse strategies, such as repetition and semantic encoding that relies significantly on the executive component. Regardless of protocol intensity and length, significant improvement in WM was seen, measured by digit span tasks.

Computerised programmes were used in six studies, all with children older than 7 years of age: Two large-scale studies (Butler et al., 2008; Butler & Copland, 2002) tested the Cognitive Remediation Program (CRP) and four studies (Bangirana et al., 2011, 2009; Boivin et al., 2010; Hardy, Willard, & Bonner, 2011) used Captain's Log computerized cognitive rehabilitation therapy (CCRT) (Sandford, 2007). In addition, one study used the Junior Cogmed Working Memory Training (JCWMT) on 21 children aged 7-12years who had Down syndrome (Bennett, Holmes, & Buckley, 2013). Both CRP and CCRT rely heavily on attention practice as a major part of the intervention, whereas JCWMT mainly consists of WM tasks that are embedded within a game-style environment. In addition, it is important to note that CCRT is not a memory- and attention-specific programme. WM improvement was seen using both longer (more than 12 rehabilitation meetings) and more intense intervention protocols (Bangirana et al., 2009; Butler et al., 2008; Butler & Copland, 2002; Hardy et al., 2011) as well as using shorter protocols, e.g., up to 10 sessions or fewer (Boivin et al., 2010). Finally, earlier intervention, three months post-injury yielded no WM improvement in comparison to later interventions (Bangirana et al., 2011).

In addition to comprehensive and computerised strategic protocols, one study (Loomes et al., 2008) examined the improvement of WM using silent rehearsal among a younger population (children aged 4–11 years). According to this study, children who rehearsed demonstrated WM improvement that was validated using a digit span task. No information was supplied on generalisation in this study.

Six studies examined semantic memory (SM) improvement after a comprehensive memory strategy approach (Patel et al., 2009; Sjö et al., 2010; van't Hooft et al., 2005, 2007, 2003; van't Hooft & Norberg, 2009). Apparently longer and more intense rehabilitation protocols (van't Hooft et al., 2005, 2007, 2003; van't Hooft & Norberg, 2009) yielded more significant improvement in SM in comparison to shorter and less intense rehabilitation protocols (Patel et al., 2009; Sjö et al., 2010). Similar results emerge when testing the efficacy of computerised programmes (CRP and CCRT). Apparently, the use of a short intervention protocol did not improve SM (Boivin et al., 2010) as compared with longer and more intense rehabilitation protocols (4–6 times of practice per week, more than 12 rehabilitation meetings) (Bangirana et al., 2009; Butler et al., 2008; Butler & Copland, 2002; Hardy et al., 2011).

In addition to comprehensive strategic protocols, several studies examined the effect of specific memory strategies, such as silent rehearsal, semantic clustering, retrieval practice and elaborative encoding on SM (Conners, Rosenquist, Arnett, Moore, & Hume, 2008; Coyne, Borg, DeLuca, Glass, & Sumowski, in press; King, White, McKinstry, Noetzel, & DeBaun, 2007; Loomes et al., 2008; Oberg & Turkstra, 1998; Yerys et al., 2003). Overall, specific memory strategies seem to be effective in children. Improvements in simple SM tasks, such as remembering simple words, were seen in all studies (Conners et al., 2008; Coyne et al., in press; Loomes et al., 2008), whereas improvement in more complex SM tasks, such as sentence memory, were seen in longer intervention protocols (King et al., 2007; Oberg & Turkstra, 1998; Yerys et al., 2003).

Errorless learning. Five studies tested the feasibility of errorless learning on SM of children; three of which are case studies. A study of two amnesic children (3 and 6 years at injury; 12 and 11 years of age at test, respectively) (Guillery-Girard, Martins, Parisot-Carbuccia, & Eustache, 2004) found that, although at a slower rate than controls, these children were still able to acquire new concepts using errorless learning approaches, despite profound episodic memory difficulties. Using the same materials and paradigm, Guilerry-Girard and colleagues (2004) reported two further cases of early onset amnesia (6 and 7 years at injury; 9.5 and 7 years of age at test, respectively) with similar findings. However in this case, only the patient with some preservation of episodic memory acquired new concepts at a rate comparable to controls. In the second case study (Martins, Guillery-Girard, Jambaqué, Dulac, & Eustache, 2006) conducted at the same centre, learning was more efficient under errorless conditions for a 15-year-old boy, although access to the information from long-term memory remained cue dependent (Martins et al., 2006).

On a larger scale study, Pauly-Takacs (2012) compared errorless learning with trial-and-error learning of declarative facts in 34 children (aged 6-16 years) with memory disorders secondary to TBI. The results did not support the use of errorless learning as a generalised intervention for learning difficulties that occur after TBI. However, in a recent study, the principle of errorless learning was more effective than trial-and-error learning in helping 15 young people with ABI (Landis et al., 2006). These inconsistent results call for extended research on the effectiveness of incorporating implicit learning techniques, such as use of errorless learning, in memory rehabilitation programmes for children.

Implicit memory

Compensation: External devices. Studies that utilised external memory aids were mainly concerned with PM. A series of studies found promising evidence for diary use in the AMAT-C protocol and in other memory rehabilitation programmes (Patel et al., 2009; van't Hooft et al., 2005, 2007, 2003). These studies demonstrated a major improvement in PM functioning in children aged 9-16 years.

In addition to diaries, the use of electronic devices was also suggested as an effective means for improving PM. For example, Ho et al. (2011) demonstrated the effectiveness of NeuroPage, a paging system in which participants were sent reminder messages regarding the tasks they needed to complete. Other electronic devices, such as personal digital assistance systems, were also found to be effective in another study with a relatively small sample of children and adolescents from a different centre (N = 12, 8–17 years) (Wilson et al., 2009). It is important to note that in the long-term, participants demonstrated spontaneous use of the pager to carry out diverse activities and used it freely, pointing to the potential efficacy of use of such a compensatory means in improving PM performance.

Behaviour management. Three studies investigated the influence of behaviour management through reinforcement learning on PM deficits by examining the role of monetary incentives in remediation following paediatric TBI. Results indicated that the high motivation (dollar per point) condition was more effective than the low motivation (penny per point) condition. However, performance of children with severe TBI in the high motivation condition was still significantly lower than controls with orthopaedic injury, and those with mild TBI in both the high and low motivation conditions (DePompei et al., 2008). This suggests that reinforcement may support the efficacy of compensatory means, but it may not necessarily mediate the deficit itself. Another type of behaviour management, conducted through parental mediation, was used in one study and demonstrated promising results on explicit visual memory in young children (aged 1.4–5 years) (McCauley & Pedroza, 2010; McCauley et al., 2009).

In addition, two studies examined how the inclusion of psycho-educational and parental support, either with or without implementing strategies to improve memory performance, supports memory performance in children with CNS disorders. Both studies found that after training/supplying information and coaching, stress levels were lower for both parents and children, and that there was an improvement in explicit WM and SM functioning (Boivin et al., 2013; van't Hooft & Norberg, 2009). This suggests the efficacy of parental support in mediating memory improvement in children with memory deficits.

Generalisation and memory systems. Generalisation of intervention effects is often challenging and is under-studied, limiting the ability to evaluate it. Nevertheless, it was found that short-term generalisation was apparent in seven studies focusing on explicit SM and WM (Butler et al., 2008; King et al., 2007; Oberg & Turkstra, 1998; van't Hooft et al., 2005; van't Hooft et al., 2007; van't Hooft et al., 2003; van't Hooft & Norberg, 2009). This is compatible with the notion that generalisation is one of the attainable goals of memory interventions for paediatric cohorts. Of these studies, five used comprehensive, long and intensive protocols, i.e., more than 12 meetings, usually several times a week (such as in the AMAT-C or CRP interventions programme), indicating, to some extent, near generalisation as well. The effect was especially seen in executive control abilities such as problemsolving skills that were not part of the original targets of the intervention. As for implicit memory, in all of the studies that used compensation aids (e.g., a diary or a smartphone) short-term generalisation was evident (DePompei et al., 2008; Ho et al., 2011; Wilson et al., 2009), such that participants were able to use the reminders in tasks other than the primary task that was tested.

DISCUSSION

Overall, the present review examined the efficacy of memory rehabilitation methods in paediatric cohorts, as a function of the target memory system and rehabilitation method. The review reveals that, especially in the last decade, remedial memory plans for children are used in diverse neurological

populations, and in multiple centres around the world and that there is a growing interest in the field of paediatric memory rehabilitation.

The responsiveness of children's discrete memory systems to rehabilitation

The first aim of this review was to investigate the responsiveness of memory systems in children. Findings showed that (1) most of the work in the area focused on WM and SM; (2) the majority of research focuses on children aged 6-17; (3) most of these studies used restoration methods; and (4) some generalisation was noted, mainly for explicit WM and implicit PM. The fact that remedial methods yielded significant improvement, along with the finding that these restorative plans demonstrate some generalisation post-treatment for WM, is consistent with previous findings (Ponsford et al., 2001; van't Hooft & Norberg, 2009) and may resonate with the notion that the WM system benefits more from increased plastic processes during late childhood.

In this context, it is interesting to compare memory system responsiveness in children to that of adults in order to better understand the factors that may promote paediatric memory rehabilitation. In the adult literature, a moderate ability to learn strategies that improve SM is reported to be helpful in real-life contexts outside the treatment context, and near generalisation is seen in patients with mild but not severe memory impairments (Rabipour & Raz, 2012). These findings are somewhat different from the studies exploring paediatric cohorts, in which even children with moderate to severe memory impairments demonstrated memory improvement in response to diverse restoration methods (Bergquist et al., 2009; Glisky & Schacter, 1986; Hampstead et al., 2012; Miller, 1992; Sohlberg & Mateer, 2001). In addition, in many cases with children, near generalisation was also seen. This difference between children and adults is noteworthy, suggesting that brain plasticity may be responsible for this finding in children.

The literature debates the possible benefits of brain plasticity in clinical paediatric populations (Boivin et al., 2010; Butler & Copland, 2002; Conners, 2003; Coyne et al., in press; King et al., 2007; Loomes et al., 2008; Oberg & Turkstra, 1998; Rankin & Hood, 2005). While increased neuroplasticity of the developing brain has generally been thought of as beneficial, it is also known that abnormal neural connectivity can result in impaired functioning and is thought of as "bad plasticity" (Giza & Prins, 2006). Nevertheless, this plasticity debate usually concerns younger children, less than three years old (Wass et al., 2012). A recent review on the effect of laboratory-based training and neurobiologically ecological interventions highlights the importance of conducting school and family-centred prevention interventions in order to improve outcomes for young (4–6 years of

age) high-risk children. The authors proposed that the influence of neurobiologically ecological interventions should be explored, especially their impact on executive function neural plasticity (Anderson et al., 2010, 2009; Giza & Prins, 2006; Montour-Proulx et al., 2004; Wass et al., 2012). This notion resonates well with another finding in this review. Indeed, we found that most of the studies examining the outcome of memory rehabilitation were conducted with children older than seven years, with little focus on younger populations. It is not known why there is a literature gap with the younger ages. Such a gap may theoretically arise from more challenges in administering the interventions at younger ages, or may indicate the need to tailor interventions that are better geared to the young, who have immature memory systems and immature executive functions. This direction of thought fits with the finding of a sharp increase in memory performance up to eight years of age, after which performance levels off (Bryck & Fisher, 2012). The use of memory rehabilitation techniques that rely on executive components and focus mainly on WM (Gathercole, 1999; Isaacs & Vargha-Khadem, 2011) after seven years of age may yield better outcomes (Bryck & Fisher, 2012; Rabipour & Raz, 2012). More specifically, it seems plausible to utilise the continued development of top-down enhancement and inhibitory control processes, which are developing in middle childhood, in rehabilitation programmes (Anderson, Anderson, Northam, Jacobs, & Catroppa, 2001; Brocki & Bohlin, 2004). This may highlight the possible mediating roles of executive functions and metacognitive abilities in supporting memory, and may indicate uncertainty regarding whether intervention efficacy is actually due to an improvement in the core memory deficit itself, or whether it is accelerated due to the development of executive functions that occurs at this age. In any event, the mediating roles of executive functions in childhood memory rehabilitation may call for structuring teaching methods to fit appropriately within the individual's functional ability. This may contribute to improvement in explicit memory performance, and brain plasticity may be increased. Future studies with fMRI or DTI that target prefrontal networks may provide further support for this notion.

In addition, an ecological factor may play an important role in the good efficacy after seven years of age. Children at school age have many realworld opportunities to practise memory techniques, especially when learning new materials in school. In contrast, for adults there are typically fewer realworld opportunities to employ the newly learned techniques. For example, most adults with memory-related pathologies are less expected to learn new lists. This suggests that the primary school age range is a target period that may benefit from specific remedial intervention, since "good plasticity" capacity and real-world opportunities are aligned and may interact to improve memory, even in children with severe memory deficits.

In this context it is important to note that lack of evidence related to younger children does not equate to lack of efficacy, but rather may denote a gap in the literature. More research is needed to understand the efficacy of memory rehabilitation programmes on younger children.

The responsiveness of children's discrete memory systems to a specific rehabilitation approach

The second aim was to explore the effectiveness of cognitive rehabilitation techniques for children. Restorative methods employ the teaching of specific memory restoration strategies, often in order to improve explicit SM and WM in children, along with generalisation effects, especially in WM practice. Additionally, the use of comprehensive programmes and a strategy based on diverse memory rehabilitation programmes have shown themselves to be effective. These data support the notion that explicit WM is easier to manipulate and improvements are noted in response to the use of multiple strategies.

More specifically, comprehensive one-on-one strategic memory rehabilitation protocols, such as AMAT-C, seem to produce beneficial effects on explicit SM and WM performance and enable generalisation in children; while use of computerised programmes produce only modest beneficial effects. This discrepancy may be due to several differences between comprehensive oneon-one rehabilitation programmes and computerised programmes. Intervention length is a possible confounding variable that may account for some of the differences between the two intervention types, because the computerised programmes are implemented for less time than the one-on-one restorative programmes (Table 1). This may indicate that restorative methods can become more effective when practised intensively for a prolonged period of time (Sander, Werkle-Bergner, Gerjets, Shing, & Lindenberger, 2012). Other reasons for differences between intervention types are related to the importance of socio-emotional support and reflective processes. The connection between emotional factors and learning was previously documented (Barnes, 1999). In terms of intervention, the child's psychosocial environment possesses necessary stability and resources to support rehabilitation goals (Laukenmann et al., 2003). Although most of the computerised and one-on-one programmes took place in the child's home environment, reflective methods are typically not used in the computerised programmes (Limond, Adlam, & Cormack, 2014). In contrast, the one-on-one programmes also included a weekly meeting with a therapist in a hospital or clinic. The purpose of these meetings is to give feedback and reinforcement, enabling the child to share her/his cognitive, emotional, and behavioural experiences. In so doing, the personal meeting provides emotional support for the child and helps maintain motivation that may, in turn, increase efficacy.

In addition, monitoring the child's progress using reflective methods may help reinforce the generalisation processes in the one-on-one programmes by strengthening meta-memory capacities and supporting internal motivation processes (Bangirana et al., 2011, 2009). This may speak to the importance of treating emotional factors as seriously as memory interventions in rehabilitation programmes because of the major influence that emotional factors have on memory and learning. Indeed, this conclusion has been applied to some comprehensive memory interventions that consider emotional support to be a critical part of the protocol (van't Hooft et al., 2005), as well as in a recent model for paediatric neuro-rehabilitation (Patel et al., 2009; Sjö et al., 2010; van't Hooft et al., 2005, 2007, 2003; van't Hooft & Norberg, 2009). More specifically, Limond and colleagues (2014) recently suggested that the following key clinical areas should be considered in paediatric rehabilitation: pragmatic factors and environmental adaptations, family functioning, challenging behaviour, emotional competence (e.g., a child's ability to recognise emotion, theory of mind), mental health of the child and the family, and motivational factors. These factors provide the foundation for effective paediatric neuro-rehabilitation tests.

In the context of providing emotional support, it seems worth considering reward and the role of behaviour management in memory rehabilitation. The effectiveness of incorporating reinforcements (Limond et al., 2014), psychoeducational support (McCauley et al., 2009; McCauley & Pedroza, 2010) and parental mediation (Ponsford et al., 2001; van't Hooft & Norberg, 2009) on memory deficits in children with CNS disorders was reported in the current review. This underscores the importance of incorporating behaviour management methods, both in the sense of direct intervention (such as reinforcement) or indirect intervention (supporting and focusing on parents), in rehabilitation programmes for children. Furthermore, the study dealing with young children (less than 5 years old) demonstrated that at this age, rehabilitation programmes rely heavily on parental mediation and support (Boivin et al., 2013). Nevertheless, only two studies investigated the efficacy of memory interventions that relied directly on parental support or parent-specific resources (Boivin et al., 2013; van't Hooft & Norberg, 2009). PM, which relies heavily on external monitoring, may benefit from interventions that combine family support with traditional rehabilitation. In addition, given the relationship between negative life events, parental psychological symptoms, and behaviour problems in children with chronic health conditions (Boivin et al., 2013; van't Hooft & Norberg, 2009), research dealing with both direct and indirect effects of family and peer support in the implementation of specific memory interventions (for both PM and explicit memory) is warranted.

In comparison with restorative methods, compensation devices that rely heavily on procedural memory may highlight the supremacy of compensation devices in improving PM. In this analysis, we found that the efficiency of using compensation devices was high even when they were used to treat children with moderate to severe memory deficits. Thus, the incorporation of memory methods that are based on procedural learning may have a beneficial effect in the paediatric population, especially when applied to everyday tasks.

Interestingly, these results are similar to those seen regarding PM improvement in adults. In one adult study, for example, compensation devices were incorporated in PM rehabilitation with promising results (Drotar, 1997; Waaland & Raines, 1991). This similarity may be explained using a developmental perspective. According to Fish et al. (2010) and Reber (1992), PM develops early in life, thus the difference between children and adults is small in their ability to rely on PM when using compensation methods during rehabilitation.

The assimilation of using compensation devices and newly learned strategic methods is partly dependent upon a psycho-education framework and on metacognitive abilities. In this review, we discovered that psycho-education has only been studied directly so far in the context of SM. Nevertheless, the results point to the important role of meta-memory, mediated by authoritative figures and significant carers, in reducing anxiety, optimising early management, and reducing the attribution of pre-existing problems to injury. This may imply that psycho-education is effective because it prevents secondary and tertiary processes by providing plans, real-time support for emotional regulation, and post hoc reflection.

Generalisation

In this context it is important to refer also to early and late generalisation. Comprehensive protocols that incorporated executive function, such as AMAT-C or CRP, demonstrated some late generalisation, especially in regard to executive control abilities such as problem-solving skills.

On the other hand, early generalisation was noted in studies that focused on implicit memory using compensation aids (such as a diary or a Smartphone). This difference raises the question of what conditions contribute to late generalisation. Is it possible that implicit memory interventions result in late generalisation as well? More research is needed to answer these questions.

Under-studied domains and future directions

We noted a lack of studies focusing on group settings of peer or family members in the context of cognitive rehabilitation. All intervention methods included in this review used a one-on-one design, precluding exploration of the therapeutic effect of peer support (Sander et al., 2012). In addition, few studies systematically integrated direct parental support or other behaviour management methods in order to improve efficacy (Funck-Brentano et al., 2005; Varni, Katz, Colegrove, & Dolgin, 1993). These areas highlight a gap in research that needs to be filled to better understand memory intervention programmes for children suffering from CNS disorders.

Finally, the interaction effects of intervention type, behaviour management method and psycho-education setting may be explored further to better understand the mechanisms involved in intervention efficacy in paediatric cohorts that are neuro-pathologically compromised.

In conclusion, the current literature provides encouraging results, suggesting that interventions effectively alleviate WM, SM and PM difficulties. We found that teaching restoration strategies may improve and result in the generalisation of SM and WM performance in children older than seven years with mild to moderate memory deficits. Further, factors such as long protocols, emotional support, and personal feedback contribute to the efficacy of restoration interventions. In addition, using compensation aids seems to be highly effective in PM tasks.

This review also highlights important under-studied areas in this field, such as the need for exploration of younger participants, group settings, and behaviour management methods. Future studies that fill these gaps may provide important contributions to the evolving field of paediatric rehabilitation and help create the most effective and appropriate guidelines for both service providers and service commissioners.

REFERENCES

- Anderson, V., Anderson, P., Northam, E., Jacobs, R., & Catroppa, C. (2001). Development of executive functions through late childhood and adolescence in an Australian sample. *Developmental Neuropsychology*, 20, 385–406. doi:10.1207/S15326942DN2001_5
- Anderson, V., Jacobs, R., Spencer-Smith, M., Coleman, L., Anderson, P., Williams, J., ... Leventer, R. (2010). Does early age at brain insult predict worse outcome? Neuropsychological implications. *Journal of Pediatric Psychology*, 35, 716–727. doi:10.1093/jpepsy/jsp100
- Anderson, V., Spencer-Smith, M., Leventer, R., Coleman, L., Anderson, P., Williams, J., ... Jacobs, R. (2009). Childhood brain insult: Can age at insult help us predict outcome? *Brain*, 1332, 45–56.
- Axelrod, B., Heilbronner, R., Barth, J., Larrabee, G., Faust, D., Pliskin, N., ... Silver, C. (2002). Cognitive rehabilitation: Official statement of the National Academy of Neuropsychology. Denver, CO: National Academy of Neuropsychology.
- Baddeley, A. (1997). Human memory: Theory and practice. Hove: Psychology Press.
- Baddeley, A. (2000). The episodic buffer: a new component of working memory? *Trends in Cognitive Sciences*, 4, 417–423. doi:10.1016/S1364-6613(00)01538-2
- Baddeley, A., & Hitch, G. (1994). Developments in the concept of working memory. *Neuropsychology*, 8, 485–493. doi:10.1037/0894-4105.8.4.485
- Bangirana, P., Allebeck, P., Boivin, M., John, C., Page, C., Ehnval, A., & Musisi, S. (2011). Cognition, behaviour and academic skills after cognitive rehabilitation in Ugandan children surviving severe malaria: A randomised trial. *Neurology*, 11, 96–103.

- Bangirana, P., Giordani, B., John, C., Page, C., Opoka, R., & Boivin, M. (2009). Immediate neuropsychological and behavioral benefits of computerized cognitive rehabilitation in Ugandan pediatric cerebral malaria survivors. *Journal of Developmental & Behavioral Pediatrics*, 30, 310–318. doi:10.1097/DBP.0b013e3181b0f01b
- Barnes, M. (1999). Rehabilitation after traumatic brain injury. British Medical Bulletin, 55, 927–943. doi:10.1258/0007142991902727
- Bauer, P., DeBoer, T., & Lukowski, A. (2007). In the language of multiple memory systems, defining and describing developments in long-term explicit memory. In L. Oakes & P. Bauer (Eds.), Short- and long-term memory in infancy and early childhood: Taking the first steps towards remembering (pp. 240–270). New York: Oxford University Press.
- Belmont, J., & Butterfield, E. (1969). The relations of short-term memory to development and intelligence. In C. Lipsett & H. Reese (Eds.), Advances in child development and behavior (Vol. 4, pp. 29–82). New York: Academic Press.
- Bennett, S., Holmes, J., & Buckley, S. (2013). Computerized Memory Training Leads to sustained improvement in visuospatial short-term memory skills in children with down syndrome. American Journal On Intellectual and Developmental Disabilitties, 118, 179–192. doi:10.1352/1944-7558-118.3.179
- Bergquist, T., Gehl, C., Mandrekar, J., Lepore, S., Hanna, S., Osten, A., & Beaulieu, W. (2009). The effect of internet-based cognitive rehabilitation in persons with memory impairments after severe traumatic brain injury. *Brain Injury*, 23, 790–799. doi:10.1080/02699050903196688
- Boivin, M., Bangirana, P., Nakasujja, N., Page, C., Sohet, C., Givon, D., ... Klein, P. (2013). A Year-Long Caregiver Training Program Improves Cognition in Preschool Ugandan Children with HIV. *Journal of pediatrics*, 163, 1409–1416. doi:10.1016/j.jpeds.2013. 06.055
- Boivin, M., Busman, R., Parikh, S., Bangirana, P., Page, C., Opoka, R., & Giordani, B. (2010). A pilot study of the neuropsychological benefits of computerized cognitive rehabilitation in Ugandan children with HIV. *Neuropsychology*, 24, 667–673. doi:10.1037/a0019312
- Braga, L., Da Paz, A., & Ylvisaker, M. (2005). Direct clinician-delivered versus indirect familysupported rehabilitation of children with traumatic brain injury: a randomized controlled trial. *Brain Injury*, 19, 819–831. doi:10.1080/02699050500110165
- Brocki, K., & Bohlin, G. (2004). Executive functions in children aged 6 to 13: A dimensional and developmental study. *Developmental Neuropsychology*, 26, 571–593. doi:10.1207/ s15326942dn2602_3
- Bryck, R., & Fisher, P. (2012). Training the brain: practical applications of neural plasticity from the intersection of cognitive neuroscience, developmental psychology, and prevention science. *The American Psychologist*, 67, 87–100. doi:10.1037/a0024657
- Butler, R., & Copland, D. (2002). Attentional processes and their remediation in children treated for cancer: A literature review and the development of a therapeutic approach. *Journal of the International Neuropsychological Society*, 8, 115–124. doi:10.1017/ S1355617702811110
- Butler, R., Copeland, D., Fairclough, D., Mulhern, R., Katz, E., Kazak, E., ... Sahler, O. (2008). A multicenter, randomized clinical trial of a cognitive remediation program for childhood survivors of a pediatric malignancy. *Journal of Consulting and Clinical Psychology*, 76, 367–378. doi:10.1037/0022-006X.76.3.367
- Carew, T., & Magsamen, S. (2010). Neuroscience and education: an ideal partnership for producing evidence-based solutions to Guide 21(st) Century Learning. *Neuron*, 67, 685–688. doi:10.1016/j.neuron.2010.08.028
- Cicerone, K., Dahlberg, C., Kalmar, K., Langenbahn, D., Malec, J., Bergquist, T., ... Morse, P. (2000). Evidence-based cognitive rehabilitation: recommendations for clinical practice. *Archives* of Physical Medicine and Rehabilitation, 81, 1596–1615. doi:10.1053/apmr.2000.19240

- Cicerone, K., Dahlberg, C., Malec, J., Langenbahn, D., Felicetti, T., Kneipp, S., ... Harley, J. (2005). Evidence-based cognitive rehabilitation: Updated review of the literature from 1998 through 2002. Archives of Physical Medicine and Rehabilitation, 86, 1681–1692. doi:10.1016/j.apmr.2005.03.024
- Cicerone, K., Langenbahn, D., Braden, C., Malec, J., Kalmar, K., Fraas, M., ... Ashman, T. (2011). Evidence-Based Cognitive Rehabilitation: Updated review of the literature from 2003 through 2008. Archives of Physical Medicine and Rehabilitation, 92, 519–530. doi:10.1016/j.apmr.2010.11.015
- Conners, F. (2003). Phonological working memory difficulty and related interventions. In J. Rondal & S. Buckley (Eds.), *Speech and language intervention in Down syndrome* (pp. 31–48). London: Colin Whurr.
- Conners, F., Rosenquist, C., Arnett, L., Moore, M., & Hume, L. (2008). Improving memory span in children with Down syndrome. *Journal of Intellectual Disability Research*, *52*, 244–255. doi:10.1111/j.1365-2788.2007.01015.x
- Coyne, J., Borg, J., DeLuca, J., Glass, L., & Sumowski, J. (in press). Retrieval Practice as an Effective Memory Strategy in Children and Adolescents with Traumatic Brain Injury. *Archives of Physical Medicine and Rehabilitation*. doi:10.1016/j.apmr.2014.09.022
- DePompei, R., Gillette, Y., Goetz, E., Xenopoulos-Oddsson, A., Bryen, D., & Dowds, M. (2008). Practical applications for use of PDAs and smartphones with children and adolescents who have traumatic brain injury. *Neurorehabilitation*, 23, 487–499.
- Domjan, M. (2003). *The principles of learning and behavior: Active learning edition* (6th Ed.). Austin: University of Texas.
- Donders, J. (2007). Traumatic brain injury. In S. J. Hunter & J. Donders (Eds.), *Pediatric neuropsychological intervention* (pp. 1–110). Cambridge: Cambridge University Press.
- Drotar, D. (1997). Relating parent and family functioning to the psychological adjustment of children with chronic health conditions: What have we learned? What do we need to know? *Journal of Pediatric Psychology*, 22, 149–165. doi:10.1093/jpepsy/22.1.1
- Ewing-Cobbs, L., Fletcher, J., Levin, H., Iovino, I., & Miner, M. (1998). Academic achievement and academic placement following traumatic brain injury in children and adolescents: A two-year longitudinal study. *Journal of Clinical and Experimental Neuropsychology*, 20, 769–781. doi:10.1076/jcen.20.6.769.1109
- Feuerstein, R. (1980). Instrumental enrichment: Redevelopment of cognitive functions of retarded performers. New York: University Park Press.
- Fish, J., Wilson, B., & Manly, T. (2010). The assessment and rehabilitation of prospective memory problems in people with neurological disorders: a review. *Neuropsychological Rehabilitation*, 20, 161–179. doi:10.1080/09602010903126029
- Funck-Brentano, I., Dalban, C., Veber, F., Quartier, P., Hefez, S., Costagliola, D., & Blanche, S. (2005). Evaluation of a peer support group therapy for HIV-infected adolescents. *AIDS*, 19, 1501–1508. doi:10.1097/01.aids.0000183124.86335.0a
- Gathercole, S. (1999). Cognitive approaches to the development of short-term memory. *Trends* in Cognitive Sciences, 3, 410–419. doi:10.1016/S1364-6613(99)01388-1
- Giza, C., & Prins, M. (2006). Is being plastic fantastic? Mechanisms of altered plasticity after developmental traumatic brain injury. *Developmental Neuroscience*, 28, 364–379. doi:10. 1159/000094163
- Glisky, E., & Schacter, D. (1986). Remediation of organic memory disorders: Current status and future prospects. *The Journal of Head Trauma Rehabilitation*, 1, 54–63. doi:10.1097/ 00001199-198609000-00009
- Gorman, S., Barnes, M., Swank, P., Prasad, M., & Ewing-Cobbs, L. (2012). The effects of pediatric traumatic brain injury on verbal and visual-spatial working memory. *Journal of the International Neuropsychological Society*, 18, 29–38. doi:10.1017/S1355617711001251

- Guillery-Girard, B., Martins, S., Parisot-Carbuccia, D., & Eustache, F. (2004). Semantic acquisition in childhood amnesic syndrome: a prospective study. *Neuroreport*, 15, 377–381. doi:10.1097/00001756-200402090-00033
- Hampstead, B., Sathian, K., Phillips, P., Amaraneni, A., Delaune, W., & Stringer, A. (2012). Mnemonic strategy training improves memory for object location associations in both healthy elderly and patients with amnestic mild cognitive impairment: A randomized, single-blind study. *Neuropsychology*, 26, 385–399. doi:10.1037/a0027545
- Hardy, K., Willard, V., & Bonner, M. (2011). Computerized cognitive training in survivors of childhood cancer: A pilot study. *Journal of Pediatric Oncology Nursing*, 28, 27–33. doi:10. 1177/1043454210377178
- Haslam, C., Bazen-Peters, C., & Wright, I. (2012). Errorless learning improves memory performance in children with acquired brain injury: a controlled comparison of standard and self-generation techniques. *Neuropsychological Rehabilitation*, 22(5), 697–715.
- Hawley, C. (2004). Behaviour and school performance after brain injury. *Brain Injury*, 18, 645–659. doi:10.1080/02699050310001646189
- Hawley, C., Ward, A., Magnay, A., & Mychalkiw, W. (2004). Return to school after brain injury. Archives of Disease in Childhood, 89, 136–142. doi:10.1136/adc.2002.025577
- Herrmann, D., Brubaker, B., Yoder, C., Sheets, V., & Tio, A. (1999). Devices that remind. In F. T. Durso (Ed.), *Handbook of applied cognition* (pp. 377–407). New York: Wiley.
- Ho, J., Epps, A., Parry, L., Poole, M., & Lah, S. (2011). Rehabilitation of everyday memory deficits in paediatric brain injury: Self-instruction and diary training. *Neuropsychological Rehabilitation*, 21, 183–207. doi:10.1080/09602011.2010.547345
- Isaacs, E., & Vargha-Khadem, F. (2011). Differential course of development of spatial and verbal memory span: A normative study. *British Journal of Developmental Psychology*, 7, 377–380. doi:10.1111/j.2044-835X.1989.tb00814.x
- Jarrold, C., Baddeley, A., & Phillips, C. (1999). Down syndrome and the phonological loop: The evidence for, and importance of, a specific verbal short-term memory deficit. *Down Syndrome Research and Practice*, 6, 61–75. doi:10.3104/reviews.97
- Jarrold, C., Purser, H., & Brock, J. (2012). Short-term memory in Down syndrome. In T. P. Alloway & S. E. Gathercole (Eds.), Working memory and neurodevelopmental disorders (pp. 239–266). NewYork: Psychology Press.
- Johnson, B., & Eagly, A. (2000). Quantitative synthesis of social psychological research. In H. T. Reis & C. M. Judd (Eds.), *Handbook of research methods in social and personality psychology* (pp. 496–528). London: Cambridge University Press.
- Jolles, D., & Crone, E. (2012). Training the developing brain: a neurocognitive perspective. *Frontiers in Human Neuroscience*, 9, 76. doi:10.3389/fnhum.2012.00076
- Karch, D., Albers, L., Renner, G., Lichtenauer, N., & von Kries, R. (2013). The efficacy of cognitive training programs in children and adolescents: a meta-analysis. *Deutsches Ärzteblatt International*, 110, 643–652.
- Kerns, K.A., & Thompson, J. (1998). Implementation of a compensatory memory system in a school age child with severe memory impairment. *Pediatric Rehabilitation*, 2(2), 77–87.
- King, A., White, D., McKinstry, M., Noetzel, M., & DeBaun, M. (2007). A pilot randomized education rehabilitation trial is feasible in sickle cell and strokes. *Neurology*, 68, 2008–2011. doi:10.1212/01.wnl.0000264421.24415.16
- Kinsella, G., Prior, M., Sawyer, M., Murtagh, D., Eisenmajer, R., Anderson, V., ... Klug, G. (1995). Neuropsychological deficit and academic performance in children and adolescents following traumatic brain injury. *Journal of Pediatric Psychology*, 20, 753–767. doi:10. 1093/jpepsy/20.6.753
- Kochanek, P. (2006). Pediatric traumatic brain injury: quo vadis?. Developmental Neuroscience, 28, 244–255. doi:10.1159/000094151

- Kodituwakku, P. (2009). Neurocognitive profile in children with fetal alcohol spectrum disorders. Developmental Disabilities Research Reviews, 15, 218–224. doi:10.1002/ddrr.73
- Kolk, A., Ennok, M., Laugesaar, R., Kaldoja, M., & Talvik, T. (2011). Long-term cognitive outcomes after pediatric stroke. *Pediatric Neurology*, 44, 101–109. doi:10.1016/j. pediatrneurol.2010.08.012
- Laatsch, L., Harrington, D., Hotz, G., Marcantuono, J., Mozzoni, M., Walsh, V., & Hersey, K. (2007). An evidence-based review of cognitive and behavioral rehabilitation treatment studies in children with acquired brain injury. *Journal of Head Trauma Rehabilitation*, 22, 248–256. doi:10.1097/01.HTR.0000281841.92720.0a
- Landis, J., Henten, G., Levin, H., Li, X., Ewing-Cobbs, L., Duron, J., & High, W. (2006). Evaluation of the errorless learning technique in children with traumatic brain injury. Archives of Physical Medicine and Rehabilitation, 87, 799–805. doi:10.1016/j.apmr.2006.02.017
- Laukenmann, M., Bleicher, M., Fuss, S., Gläser-Zikuda, M., Mayring, P., & von Rhöneck, C. (2003). An investigation of the influence of emotional factors on learning in physics instruction. *International Journal of Science Education*, 25, 489–507. doi:10.1080/ 09500690210163233
- Limond, J., Adlam, A., & Cormack, M. (2014). A model for pediatric neurocognitive interventions: considering the role of development and maturation in rehabilitation planning. *The Clinical Neuropsychologist*, 28, 181–198. doi:10.1080/13854046.2013.873083
- Limond, J., & Leeke, R. (2005). Practitioner Review: Cognitive rehabilitation for children with acquired brain injury. *Journal of Child Psychology and Psychiatry*, 46, 339–352. doi:10. 1111/j.1469-7610.2004.00397.x
- Lindahl, N., & Steneby, S. (2009). Experiences of parents of children surviving brain tumour: a happy ending and a rough beginning. *European Journal of Cancer Care*, 18, 371–380. doi:10.1111/j.1365-2354.2008.00976.x
- Loomes, C., Rasmussen, C., Pei, J., Manji, S., & Andrew, G. (2008). The effect of rehearsal training on working memory span of children with fetal alcohol spectrum disorder. *Research* in Developmental Disabilities, 29, 113–124. doi:10.1016/j.ridd.2007.01.001
- Manji, S., Pei, J., Loomes, C., & Rasmussen, C. (2009). A review of the verbal and visual memory impairments in children with foetal alcohol spectrum disorders. *Developmental Neurorehabilitation*, 12, 239–247. doi:10.1080/17518420902980118
- Martins, S., Guillery-Girard, B., Jambaqué, I., Dulac, O., & Eustache, F. (2006). How children suffering severe amnesic syndrome acquire new concepts?. *Neuropsychologia*, 44, 2792– 2805. doi:10.1016/j.neuropsychologia.2006.05.022
- Mattson, S., & Riley, E. (2006). A review of the neurobehavioral deficits in children with fetal alcohol syndrome or prenatal exposure to alcohol. *Alcoholism: Clinical and Experimental Research*, 22, 279–294. doi:10.1111/j.1530-0277.1998.tb03651.x
- McCauley, S., & Levin, H. (2004). Prospective memory in pediatric traumatic brain injury: A preliminary study. *Developmental Neuropsychology*, 25, 5–20. doi:10.1080/87565641. 2004.9651919
- McCauley, S., McDaniel, M., Pedroza, C., Chapman, S., & Levin, H. (2009). Incentive effects on event-based prospective memory performance in children and adolescents with traumatic brain injury. *Neuropsychology*, 23, 201–209. doi:10.1037/a0014192
- McCauley, S., & Pedroza, C. (2010). Event-based prospective memory in children with sickle cell disease: Effect of cue distinctiveness. *Child Neuropsychology*, *16*, 293–312. doi:10. 1080/09297041003601470
- McCauley, S.R., Pedroza, C., Chapman, S.B., Cook, L.G., Vásquez, A.C., & Levin, H.S. (2011). Monetary incentive effects on event-based prospective memory three months after traumatic brain injury in children. *Journal of Clinical and Experimental Neuropsychology*, 33(6), 639–646. doi:10.1080/13803395.2010.547844

- McDaniel, M., & Einstein, G. (2007). Prospective memory: An overview and synthesis of an emerging field. Los Angeles, CA: Sage Publications.
- McKeough, A., Lupart, J., & Marini, A. (1995). Teaching for Transfer: Fostering Generalization in Learning. New Jersey: Lawrence Erlbaum Associates, Inc.
- Melby-Lervåg, M., & Hulme, C. (2013). Is working memory training effective?. A meta-analytic review. Developmental Psychology, 49, 270–291.
- Michaud, L., Rivara, F., Grady, M., & Reay, D. (1992). Predictors of survival and severity of disability after severe brain injury in children. *Neurosurgery*, 31, 254–264. doi:10.1227/ 00006123-199208000-00010
- Miller, E. (1992). Psychological approaches to the management of memory impairments. The British Journal of Psychiatry, 160, 1–6. doi:10.1192/bjp.160.1.1
- Montour-Proulx, L., Braun, C., Daigneault, S., Rouleau, L., Kuehn, S., & Begin, J. (2004). Predictors of intellectual function after a unilateral cortical lesion: Study of 635 patients from infancy to adulthood. *Journal of Child Neurology*, 19, 935–943.
- Mulhern, R., & Butler, R. (2004). Review Neurocognitive sequelae of childhood cancers and their treatment. *Developmental Neurorehabilitation*, 7, 1–14. doi:10.1080/ 13638490310001655528
- Mulhern, R., & Butler, R. (2006). Neuropsychological late effects. In R. Brown (Ed.), Comprehensive handbook of childhood cancer and sickle cell disease: a biopsychosocial approach (pp. 262–278). New York: Oxford University Press.
- Oberg, L., & Turkstra, L. (1998). Use of elaborative encoding to facilitate verbal learning after adolescent traumatic brain injury. *Journal of Head Trauma Rehabilitation*, 13, 44–62. doi:10.1097/00001199-199806000-00005
- Patel, S., Katz, E., Richardson, R., Rimmer, M., & Kilian, S. (2009). Cognitive and problem solving training in children with cancer: A pilot project. *Journal of Pediatric Hematology* and Oncology, 31, 670–677. doi:10.1097/MPH.0b013e3181b25a1d
- Pauly-Takacs, K. (2012). Learning and episodic memory following childhood brain tumour: a case study. Leeds West Yorkshire: University of Leeds.
- Ponsford, J., Willmot, C., Rothwell, A., Cameron, P., Ayton, G., Nelms, R., ... Ng, K. (2001). Impact of early intervention on outcome after mild traumatic brain injury. *Pediatrics*, 108, 1297–1303. doi:10.1542/peds.108.6.1297
- Rabipour, S., & Raz, A. (2012). Training the brain: fact and fad in cognitive and behavioral remediation. *Brain and Cognition*, *79*, 159–179. doi:10.1016/j.bandc.2012.02.006
- Ramsden, S., & Hubbard, J. (2002). Family expressiveness and parental emotion coaching: their role in children's emotion regulation and aggression. *Journal of Abnormal Child Psychol*ogy, 30, 657–667. doi:10.1023/A:1020819915881
- Rankin, P., & Hood, J. (2005). Designing clinical interventions for children with specific memory disorders. *Pediatric Rehabilitation*, 8, 283–297.
- Rasmussen, C. (2006). Executive functioning and working memory in fetal alcohol spectrum disorder. Alcoholism: Clinical and Experimental Research, 29, 1359–1367. doi:10.1097/ 01.alc.0000175040.91007.d0
- Reber, A. (1992). The cognitive unconscious: An evolutionary perspective. *Consciousness and Cognition*, 1, 93–133. doi:10.1016/1053-8100(92)90051-B
- Reeves, C., Palmer, S., Reddick, W., Merchant, T., Buchanan, G., Gajjar, A., & Mulhern, R. (2006). Attention and memory functioning among pediatric patients with medulloblastoma. *Journal of Pediatric Psychology*, 31, 272–280. doi:10.1093/jpepsy/jsj019
- Ross, K., Dorris, L., & McMillan, T. (2011). A systematic review of psychological interventions to alleviate cognitive and psychosocial problems in children with acquired brain injury. *Developmental Medicine and Child Neurology*, 53, 692–701. doi:10.1111/j.1469-8749. 2011.03976.x

- Sander, M., Werkle-Bergner, M., Gerjets, P., Shing, Y., & Lindenberger, U. (2012). The twocomponent model of memory development, and its potential implications for educational settings. *Developmental Cognitive Neuroscience*, 2(Suppl 1), S67–S77. doi:10.1016/j. dcn.2011.11.005
- Sandford, J. A. (2007). *Captain's log computerized cognitive training system*. Richmond, VA: Brain Train.
- Schacter, D. (1987). Implicit memory: History and current status. Journal of Experimental Psychology: Learning, Memory and Cognition, 13, 501–518. doi:10.1037/0278-7393.13.3.501

Schacter, D., & Tulving, E. (1994). What are the memory systems of 1994.

- Serra-Grabulosa, J., Junque, C., Verger, K., Salgado-Pineda, P., Maneru, C., & Mercader, J. (2005). Cerebral correlates of declarative memory dysfunctions in early traumatic brain injury. *Journal of Neurology, Neurosurgery & Psychiatry*, 76, 129–131. doi:10.1136/ jnnp.2004.027631
- Shadmehr, R., & Moussavi, Z. (2000). Spatial Generalization from Learning Dynamics of Reaching Movements. *The Journal of Neuroscience*, 20, 7807–7815.
- Sjö, M., Spellerberg, S., Weidner, S., & Kihlgren, M. (2010). Training of attention and memory deficits in children with acquired brain injury. *Acta pediatrica*, 99, 230–236. doi:10.1111/j. 1651-2227.2009.01587.x
- Slomine, B., & Locascio, G. (2009). Cognitive rehabilitation for children with acquired brain injury. *Developmental Disabilities Research Review*, 15, 133–143. doi:10.1002/ddrr.56
- Smith, L., Adnams, C., & Eley, B. (2008). Neurological and neurocognitive function of HIVinfected children commenced on antiretroviral therapy. *South African Journal of Child Health*, 2, 108–113.
- Sohlberg, M., & Mateer, C. (2001). Cognitive rehabilitation: An integrative neuropsychological approach. New York: Guilford Press.
- Souza, L., Braga, L., Filho, G., & Dellatolas, G. (2007). Quality-of-life: Child and parent perspectives following severe traumatic brain injury. *Developmental Neurorehabilitation*, 10, 35–47. doi:10.1080/13638490600822239
- Squire, L. (1992). Encyclopedia of learning and memory. New York: Macmillan Publishing Co, Inc.
- Squire, L., & Schacter, D. (2002). Neuropsychology of memory 3th edition. New York: Guilford press.
- Tatla, S., Sauve, K., Jarus, T., Virji-Babul, N., & Holsti, L. (2014). The effects of motivating interventions on rehabilitation outcomes in children and youth with acquired brain injuries: a systematic review. *Brain Injury*, 28, 1022–1035. doi:10.3109/02699052.2014.890747
- Tulving, E. (1972). Episodic and semantic memory. In E. Tulving & W. Donaldson (Eds.), Organization of memory (pp. 381–403). New York: Academic Press.
- Tulving, E., & Schacter, D. (1990). Priming and human memory systems. Science, 247, 301– 306. doi:10.1126/science.2296719
- Vakil, E. (2005). The Effect of Moderate to Severe Traumatic Brain Injury (TBI) on Different Aspects of Memory: A Selective Review. *Journal of Clinical and Experimental Neuropsychology*, 27, 977–1021. doi:10.1080/13803390490919245
- Van Heugten, C., Hendriksen, J., Rasquin, S., Dijcks, B., Jaeken, D., & Vles, J. (2006). Longterm neuropsychological performance in a cohort of children and adolescents after severe paediatric traumatic brain injury. *Brain Injury*, 20, 895–903. doi:10.1080/ 02699050600832015
- van't Hooft, I., Andersson, K., Bergman, B., Sejersen, T., von Wendt, L., & Bartfai, A. (2005). Beneficial effect from a cognitive training programme on children with acquired brain injuries demonstrated in a controlled study. *Brain Injury*, 19, 511–518. doi:10.1080/ 02699050400025224

- van't Hooft, I., Andersson, K., Bergman, B., Sejersena, T., von Wendt, L., & Bartfai, A. (2007). Sustained favorable effects of cognitive training in children with acquired brain injuries. *Neurorehabilitation*, 22, 109–116.
- van't Hooft, I., Andersson, K., Sejersen, T., Bartfai, A., & von Wendt, L. (2003). Attention and memory training in children with acquired brain injuries. *Acta pediatrica*, 92, 935–940. doi:10.1111/j.1651-2227.2003.tb00627.x
- van't Hooft, I., & Norberg, A. (2009). SMART cognitive training combined with a parental coaching programme for three children treated for medulloblastoma. *Neurorehabilitation*, 25, 1–9.
- Vargha-Khadem, F., Gadian, D., Watkins, K., Connelly, K., Van Paesschen, W., & Mishkin, M. (1997). Differential Effects of Early Hippocampal Pathology on Episodic and Semantic Memory. *Science*, 277, 376–380. doi:10.1126/science.277.5324.376
- Varni, J., Katz, E., Colegrove, R., & Dolgin, M. (1993). The impact of social skills and training on the adjustment of children with newly diagnosed cancer. *Journal of Pediatric Psychol*ogy, 18, 751–767. doi:10.1093/jpepsy/18.6.751
- Waaland, P., & Raines, S. (1991). Families coping with childhood neurological disability: Clinical assessment and treatment. *Neuropsychological Rehabilitation*, 1, 19–27.
- Ward, H., Shum, D., McKinlay, L., Baker, S., & Wallace, G. (2007). Prospective Memory and Pediatric Traumatic Brain Injury: Effects of Cognitive Demand. *Child Neuropsychology: A Journal on Normal and Abnormal Development in Childhood and Adolescence*, 13, 219–239.
- Ward, H., Shum, D., Wallace, G., & Boon, J. (2002). Pediatric traumatic brain injury and procedural memory. *Journal of Clinical and Experimental Neuropsychology*, 24, 458–470. doi:10.1076/jcen.24.4.458.1032
- Wass, S., Scerif, G., & Johnson, M. (2012). Training attentional control and working memory Is younger, better? *Developmental Review*, 32, 360–387. doi:10.1016/j.dr.2012.07.001
- Willingham, D., Nissen, M., & Bullemer, P. (1989). On the development of procedural knowledge. Journal of Experimental Psychology: Learning, Memory, and Cognition, 15, 1047– 1060. doi:10.1037/0278-7393.15.6.1047
- Wilson, B., Baddeley, A., Evans, J., & Shiel, A. (1994). Errorless learning in the rehabilitation of memory impaired people. *Neuropsychological Rehabilitation*, 4, 307–326. doi:10.1080/ 09602019408401463
- Wilson, B., Emslie, H., Evans, J., Quirk, K., Watson, P., & Fish, J. (2009). The NeuroPage system for children and adolescents with neurological deficits. *Developmental Neurorehabilitation*, 12, 421–426. doi:10.3109/17518420903200573
- Wilson, B., & Glisky, E. (2009). Memory rehabilitation: integrating theory and practice. New York: Guilford Press.
- Wilson, B., & Moffat, N. (1992). *Clinical managment of memory problems*. London: Chapman and Hall.
- Yeates, K., & Enrile, B. (2005). Implicit and explicit memory in children with congenital and acquired brain disorder. *Neuropsychology*, 19, 618–628. doi:10.1037/0894-4105.19.5.618
- Yerys, B., White, D., Salorio, C., McKinstry, R., Moinuddin, A., & DeBaun, M. (2003). Memory Strategy Training in Children With Cerebral Infarcts Related to Sickle Cell Disease. *Journal of Pediatric Hematology/Oncology*, 25, 495–498. doi:10.1097/ 00043426-200306000-00014
- Ylvisker, M., Andelson, D., Braga, L., Bumett, S., Glang, A., Feeney, T., ... Todis, B. (2005). Rehabilitation and ongoing support after paediatric brain injury: Twenty years of progress. *Journal of Head Trauma Rehabilitation*, 20, 95–109. doi:10.1097/00001199-200501000-00009

APPENDIX 1

Methodological quality appraisals of papers.

Score 1 if met, 0 if not met or unable to determine

- (1) Were specific hypotheses and or objectives stated?
- (2) Were the settings and locations where data were collected stated?
- (3) Was a control or comparison group used?
- (4) Were participants randomly allocated to groups?
- (5) Is the method of randomisation appropriate?
- (6) Was the total sample size > 20 participants?
- (7) Was the total sample size > 40 participants?
- (8) Were at least some of the measures standardised assessment tools?
- (9) Were the measures appropriate for age group?
- (10) Were the inclusion exclusion criteria clearly stated?
- (11) Did the article specify the severity of the brain injury for participants with acquired brain injury and was the method of diagnosis appropriate (e.g., by a medical professional, Glasgow Coma Scale)?
- (12) Did the injury occur at least 6 months previously (to ensure the results were not a reflection of the recovery process)?
- (13) Were follow-up data collected after post-intervention data (i.e., to see if effects were maintained post-intervention)?
- (14) Were all participants included in the analysis?
- (15) If not, was intent-to-treat analysis used? (Award 1 point if a point is granted on the above item.)
- (16) Were those assessing the outcomes blind to the group?
- (17) Was a power calculation used or sample size justified?
- (18) Was the intervention described in detail (i.e., how it was administered, etc.) or was there reference to a manual?
- (19) Were the characteristics of participants clearly described (e.g., demographic information such as age, sex)?
- (20) Did the results relate to the initial hypotheses?
- (21) Was the statistical analysis appropriate?
- (22) Were data adequately described (mean, range, etc.)?
- (23) Were effect sizes calculated?
- (24) Were effect sizes moderate or better (for studies with small sample sizes n < 10)?
- (25) Was there sufficient information to calculate effect size (i.e. mean and SD)?
- (26) Was age taken into account as a possible confounding factor?

Total quality ratings.