

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/330818208>

Inhibitory effect of rosemary essential oil, loaded in liposomes, on seed germination of *Acacia saligna* (Labill.) Wendl., an invasive species in Mediterranean ecosystems

Article in *Botany* · February 2019

DOI: 10.1139/cjb-2018-0212

CITATIONS

2

READS

304

7 authors, including:



Alfredo Maccioni

12 PUBLICATIONS 24 CITATIONS

[SEE PROFILE](#)



Andrea Santo

87 PUBLICATIONS 443 CITATIONS

[SEE PROFILE](#)



Falconieri Danilo

Istituto Tecnico Industriale M. Giua, Cagliari, Italy

84 PUBLICATIONS 1,443 CITATIONS

[SEE PROFILE](#)



Alessandra Piras

Università degli studi di Cagliari

148 PUBLICATIONS 2,825 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Distribution Patterns and Conservation Status Assessment of Exclusive Endemic Vascular Plant Taxa in Matrouh Governorate, Egypt. [View project](#)



IUCN Red List [View project](#)

Inhibitory effect of rosemary essential oil, loaded in liposomes, on seed germination of *Acacia saligna*, an invasive species in Mediterranean ecosystems

Alfredo Maccioni, Andrea Santo, Danilo Falconieri, Alessandra Piras, Maria Manconi, Andrea Maxia, and Gianluigi Bacchetta

Abstract: *Acacia saligna* (Labill.) Wendl. is native of southwestern Australia, but has been planted extensively in many areas of the world, including the Mediterranean region, becoming highly invasive especially in coastal habitats. The aim of this study was to test whether the indigenous *Rosmarinus officinalis* L. essential oil (EO), loaded in liposomes, can inhibit the seed germination of the invasive alien *Acacia saligna*. Variability in seed germination requirements and responses among populations were evaluated. Germination tests under light, at constant temperatures and three concentrations of rosemary EO were carried out. Among the examined factors, only the EO amount and temperature had a highly significant effect on seed germination. The lowest EO quantity did not show differences compared with the control, while the highest amount inhibited significantly seed germination of all populations at all the tested temperatures. No seed recovered the ability to germinate after the EO treatment with either of the two washing methods. Our results allowed us to identify the minimum amount of rosemary EO capable of inhibiting the seed germination of the invasive *Acacia saligna*. These results could be useful for the control of this invasive alien species, thus allowing the conservation of indigenous Mediterranean plant species and habitats.

Key words: essential oil, invasive alien species, liposomes, *Rosmarinus officinalis*, seed germination.

Résumé : *Acacia saligna* (Labill.) Wendl. est indigène dans le sud-ouest de l'Australie, mais elle a été largement plantée dans plusieurs régions dans le monde, y compris dans la région méditerranéenne, devenant hautement envahissante, spécialement dans les habitats côtiers. L'objectif de cette étude consistait à tester si l'huile essentielle (HE) de *Rosmarinus officinalis* L. indigène, chargée dans des liposomes, peut inhiber la germination des graines de l'espèce exotique envahissante *Acacia saligna*. La variabilité des besoins pour la germination des graines et les réponses des populations ont été évaluées. Des tests de germination sous la lumière, à des températures constantes et en présence de trois concentrations d'HE de romarin ont été réalisés. Parmi les facteurs examinés, seules la quantité d'HE et la température exerçaient un effet hautement significatif sur la germination des graines. La quantité d'HE la plus faible n'était pas différente du contrôle, alors que les quantités les plus élevées inhibaient significativement la germination des graines de toutes les populations à toutes les températures examinées. Aucune des graines ne récupérait la capacité de germer après un traitement à l'HE, quelle que soit la méthode de lavage utilisée parmi les deux qui ont été testées. Les résultats ont permis aux auteurs d'identifier une quantité minimale d'HE de romarin capable d'inhiber la germination des graines de l'espèce envahissante *Acacia saligna*. Ces résultats pourraient être utiles pour contrôler cette espèce exotique envahissante, permettant ainsi la conservation des espèces indigènes méditerranéennes et de leurs habitats. [Traduit par la Rédaction]

Mots-clés : huile essentielle, espèce exotique envahissante, liposomes, *Rosmarinus officinalis*; germination des grains.

Received 25 November 2018. Accepted 17 January 2019.

A. Maccioni and A. Maxia.* Department of Life and Environmental Sciences, University of Cagliari, viale S. Ignazio da Laconi 13, 09123, Cagliari, Italy; Co.S.Me.Se, Consorzio Interuniversitario per lo Studio dei Metaboliti Secondari, viale S. Ignazio da Laconi 13, 09123, Cagliari, Italy.
A. Santo and M. Manconi. Department of Life and Environmental Sciences, University of Cagliari, viale S. Ignazio da Laconi 13, 09123, Cagliari, Italy.
D. Falconieri. Co.S.Me.Se, Consorzio Interuniversitario per lo Studio dei Metaboliti Secondari, viale S. Ignazio da Laconi 13, 09123, Cagliari, Italy.
A. Piras. Department of Chemical and Geological Sciences, University of Cagliari, Cittadella Universitaria, S.S. 554, 09042, Monserrato (CA), Italy.
G. Bacchetta.* Department of Life and Environmental Sciences, University of Cagliari, viale S. Ignazio da Laconi 13, 09123, Cagliari, Italy; Hortus Botanicus Karalitanus (HBK), University of Cagliari, viale S. Ignazio da Laconi 13, 09123, Cagliari, Italy.

Corresponding author: Alfredo Maccioni (email: alfredomaccioni87@gmail.com).

*These authors contributed equally to the work.

Copyright remains with the author(s) or their institution(s). Permission for reuse (free in most cases) can be obtained from [RightsLink](https://www.copyright.com).

Introduction

Invasive alien species (IAS) are considered, after habitat loss and fragmentation, as one of the greatest threats to the conservation of indigenous species and habitats worldwide (Crisóstomo et al. 2013). Moreover, these species also cause significant economic damage to natural ecosystems and to human health (DAISIE 2009; Strydom et al. 2012).

IAS are responsible for the reduction of native species abundance and diversity, change in nutrient cycling, water availability, and fire regimes, altering both the structure and function of invaded ecosystems (Correia et al. 2014). Callaway and Aschehoug (2000) asserted that some IAS may be successful because they introduce novel mechanisms of interaction by releasing substances that interfere with primary and secondary physiological processes, e.g., ions and water uptake, enzyme functions, seed germination, seedling growth, photosynthesis, and respiration of native plant species (Lorenzo et al. 2011).

The genus *Acacia* Mill. belongs to the Fabaceae (nom. alt. Leguminosae) family (Richardson et al. 2011) and includes some of the most important IAS in the world. About 23 species of this genus are major invaders in Mediterranean ecosystems (Werner et al. 2010). The effects of invasion by *Acacia* species on indigenous species and habitats have been widely documented (Le Maitre et al. 2011; Correia et al. 2014), particularly on species richness, community structure, soil bulk density, and water availability (Werner et al. 2010). Moreover, these species show several reproductive traits that may also contribute to their invasiveness, such as the ability to self-pollinate, vegetative reproduction, and the production of a large amount of long-lived and highly viable seeds (Morris et al. 2011; Correia et al. 2014; Meloni et al. 2015).

Acacia saligna (Labill.) Wendl. is an evergreen phanerophyte, native of southwestern Australia (Orchard and Wilson 2001) and is considered a neophyte for the Mediterranean Basin. This species grows in a wide range of ecological conditions, including semi-arid and arid Mediterranean climatic areas, especially on barren slopes, sand dunes (Midgley and Turnbull 2003), and abandoned lands (Kutiel et al. 2004). Musil (1993) reported that *Acacia saligna* is tolerant to alkaline and saline soils, exhibiting greater tolerance to high concentrations of potassium, calcium, and magnesium compared with several indigenous species (Fox 1995; Doran and Turnbull 1997).

One of the factors that may have a strong influence on *Acacia saligna* invasiveness is its ability to produce a large number of viable seeds (Strydom et al. 2012). *Acacia saligna* seeds may remain dormant but viable for more than 50 years, until the seed coat is sufficiently damaged to be permeable to water necessary for seed imbibition (Meloni et al. 2015). According to Holmes et al. 1987 and Holmes and Cowling (1997a, 1997b) they maintain high

viability (values of ca. 86%–100%) in the soil seed bank even after several years.

During the last two decades, this species has been widely used for stabilizing shifting sand dunes, rehabilitating sandy mining areas, protecting barren rocks along roadsides, and has been planted extensively in many parts of the world, becoming highly invasive in the Mediterranean Basin (Crisóstomo et al. 2013; Del Vecchio et al. 2013).

Essential oils (EOs) are produced by more than 17 000 aromatic plant species belonging mostly to a few families, including the Myrtaceae, Lauraceae, Lamiaceae, and Asteraceae (Regnault-Roger et al. 2012). EOs contain compounds named plant secondary metabolites, such as terpenes and phenols, and are known to have multiple ecological roles, notably defence against herbivores, pathogens, and abiotic stresses (Iason et al. 2012). The production of these compounds is affected by the physical and chemical properties of the ecosystem where the plants grow, such as precipitation, temperature, and edaphic features (Thompson 2005; Pirbalouti et al. 2015). Plant secondary metabolites may be present in all plant organs, and may be found in different concentrations in several parts of the plant. The concentration of these substances is the parameter that determines a positive influence (facilitation) and (or) negative (competition) against other target organisms (Amiot et al. 2005; Iason et al. 2012). The beneficial or harmful influence of these compounds may have an influence on seed germination and seedling growth of other nearby plants, allelopathy, or phytotoxicity (Rice 1984).

The Mediterranean biogeographic region is rich in aromatic plants, many of which belong to the Lamiaceae (Chizzola 2006), which is considered one of the largest and most distinctive families of flowering plants, with about 220 genera and almost 4000 species worldwide (Mitić et al. 2011). *Rosmarinus officinalis* L. (rosemary), which has recently (2017) been placed in *Salvia* L. genus as *Salvia rosmarinus* (L.) Schleid. by Drew et al. (2017) and is known as *Salvia rosmarinus* Schleid. (Bartolucci et al. 2018), is one of the most important aromatic species of this family and it has been used since ancient times for medicinal and culinary uses (Dajic-Stevanovic et al. 2008). Rosemary is a circum-Mediterranean evergreen indigenous nanophanerophyte, calcicolous and heliophilous, which grows in many substrata, preferring sandy, dry, calcareous, and humus-poor soils. This species is characteristic of the Mediterranean maquis, garrigue and shrublands, and can be found from sea level up to 800 m a.s.l., and contains a quantity up to 1% of EO (Pintore et al. 2002; Bellumori et al. 2015).

Detailed studies of the effects of EO from several members of the Lamiaceae have shown that, depending on the type and concentration, EOs reduced or totally inhibited germination of a range of species (Angelini et al. 2003; Atak et al. 2016).

Table 1. Population data and seed lot details.

Population code	Site	Island	Mean coordinates (UTM, WGS84)	Substrate	Distance from sea (m)	Mean elevation (m a.s.l.)	Mean($\pm$ SD) seed mass (mg)
Asal1	Costa Rei, Muravera (CA)	SE Sardinia	41°14'N, 09°24'E	Aeolian sands	10	3	17.94 $\pm$ 0.54
Asal2	La Maddalena (OT)	NE Sardinia	41°35'N, 09°18'E	Granites	2	2	18.96 $\pm$ 1.04
Asal3	Arenella, Siracusa (SR)	SE Sicily	37°00'N, 15°16'E	Calcarenes	45	10	19.92 $\pm$ 0.82
Asal4	Punta del Pero, Siracusa (SR)	SE Sicily	37°02'N, 15°17'E	Limestones	24	5–8	18.37 $\pm$ 0.67

Note: CA, Cagliari; OT, Olbia-Tempio; SR, Siracusa.

The aims of this research were to obtain preliminary results on the effect of rosemary EO on *Acacia saligna* seeds, in particular: (1) to define the potential phytotoxic effect of rosemary EO on seed germination, evaluating the minimum amount of rosemary EO able to inhibit seed germination; and (2) evaluate inter-population variability in the seed germination response in *Acacia saligna*.

In this study, the EO was loaded in liposomes, which are phospholipid nanovesicles able to limit the loss of volatile components and deliver them inside the seed potentiating the EO efficacy. Additionally, the EO incorporation in liposomes ensures its complete and homogeneous dispersion in the agar reducing its loss. To our knowledge, this is the first time that liposomes have been used as components of a germination substrate.

Materials and methods

Plant material, EO, and seed lot details

Flowering aerial parts of 10 individuals of *R. officinalis* were collected in North Sardinia (Su Canale, Monti (Olbia-Tempio, Italy); 40°50'N, 9°23'E). Afterwards, the plant material was air-dried in an oven (FP 115; BINDER) at 40 °C with forced ventilation for 48 h. The plant material was subjected to steam distillation to obtain the EO. The chemical composition of EO was analysed by using gas chromatography (GC-FID) and gas chromatography – mass spectrometry (GC-MS) according to Zuzarte et al. (2013).

Acacia saligna legumes (hereinafter fruits) were collected in the period of natural dispersal (summer) in four naturalized Mediterranean populations (Table 1). Seeds were extracted from the fruits by hand and using laboratory tweezers. Prior to germination tests, seeds were stored under controlled conditions (20 °C and 50% relative humidity) for 2 weeks.

Germination trials

Germination tests were performed at the Sardinian Germplasm Bank (BG-SAR) of the Hortus Botanicus Karalitanus (HBK) of the University of Cagliari, Sardinia (Italy). To detect the presence of water-impermeable integuments, i.e., the physical component of seed dormancy, seven different methods of both mechanical and chemical scarification were tested. In each method, four replicates of 25 seeds per Petri dish were used. Furthermore, another four replicates with 25 intact seeds per Petri dish were used as the control (1). Seeds were

chipped with a scalpel (2), engraved with nail clippers (3), mechanically scarified with sandpaper No. 40 (4) and No. 80 (5), immersed in H₂SO₄ (96%) for 10 (6), 15 (7), and 20 (8) min (Azad et al. 2011). All seeds were sown under a laminar flow hood (KB; Faster) in 90 mm diameter plastic Petri dishes on 1% water agar substrate, which provided a solid, non-sterile medium for germination, independently from the treatment, and were incubated at a constant temperature of 15 °C in a growth chamber (MLR-351; SANYO) in the light (12 h of irradiance per day). After sowing, each Petri dish was sealed with laboratory parafilm (parafilm M; Pechiney) to limit evaporation and prevent contamination spores and microorganisms from the outside.

The criterion for germination was visible radicle protrusion (≥ 2 mm). Seeds were scored daily and germinated seeds discarded. When no additional germination occurred for two consecutive weeks, tests were stopped, and the viability of each remaining seed was checked by a cut test with a scalpel and subsequent observation of the endosperm under binocular microscope.

Several preliminary experimental tests were carried out to identify the more suitable formulation able to stably load the EO and easily blend with the germination substrate. In particular, during these preliminary experiments, soy lecithin, which forms liposomes, was identified as the best component for this aim.

The effect of soy lecithin (empty liposomes) on the seed germination of *Acacia saligna* was evaluated by testing four replicates with the addition of soy lecithin to the substrate and four replicates with only 1% agar. Seeds were incubated in a growth chamber at 15 °C in the light (12 h of irradiance per day). Meloni et al. (2015) showed that these conditions were optimal to achieve the highest germination for this species. For all the preliminary experiments, Asal1 seeds (Table 1) were used (due to their higher availability).

In each Petri dish used for the germination tests with the addition of EO, 50 mL of solution was used for the germination substrate. The substrate consisted of two layers: A (lower) and B (upper). Layer A contained 30 mL of solution, while the remaining 20 mL was used to create layer B on which seeds were sown. Layer A consisted of 1% agar. Layer B contained 1% agar, distilled water, and liposomes containing 1.8 g of soy lecithin and EO (on the basis of the used amount). To ensure a homogenous mix-

ing of the EO and the agar substrate, the EO was loaded in liposomes, which were prepared by sonicating the soy lecithin and the EO dispersed in water (Soniprep 150 Plus Sonicators; MSE) for 30 min, then immediately mixed with agar. It was not possible to prepare layer B without using EO-loaded liposomes because the free EO added to the agar substrate precipitated.

We tested the effect of three different concentrations of *R. officinalis* EO (3.90, 7.81, 15.62 $\mu\text{L}/50\text{ mL}$ of the substrate) loaded in liposomes containing the same amount of soy lecithin. Preliminary experiments showed that higher concentrations did not produce a solid substrate.

At the end of the experiments, to evaluate the reversibility of the EO effects, the ability of *Acacia saligna* seeds to recover their germination after the EO treatment was evaluated by washing seeds with Tween 80 PS ($\text{C}_{64}\text{H}_{124}\text{O}_{26}$, Polysorbate 80; Panreac) and sodium hypochlorite (NaClO) solutions. The Tween 80 washing lasted 10 min (the first 5 min with 200 mL/L Tween 80 in distilled water and the remaining 5 min with 100 mL/L). The sodium hypochlorite (860 mM) washing lasted 10 min (the first 5 min with 500 mL/L bleach in distilled water and the remaining 5 min with 250 mL/L). The response of *Acacia saligna* seeds from each population was evaluated on the basis of the seed availability, therefore we tested three temperatures (10, 15, 20 $^{\circ}\text{C}$) for the two Sardinian populations and two (15, 20 $^{\circ}\text{C}$) for the Sicilian ones, with photoperiod of 12 h of irradiance per day.

Data analysis

Final germination percentages were calculated as the mean of four replicates ($\pm\text{SD}$). The rate of germination was estimated by using a modified Timson's index of germination velocity (TI) (Khan and Ungar 1984; Santo et al. 2015): $\text{TI} = \sum G/t$, where G is the percentage of seed germination at 2-day interval and t is the total germination period. Arcsine-transformed germination percentages and log10-transformed TI were analysed by ANOVA and consequent Fisher's least significant differences post hoc test. Germination percentages of scarification pretreatments were analysed by the nonparametric Kruskal-Wallis test, followed by a Mann-Whitney U test, due to the non-satisfaction of the ANOVA assumptions even after arcsine transformation. Data were graphed using Sigmaplot 11.0 (Systat Software Inc., London, UK), and all statistical analyses were carried out using the statistical software Statistica 7.0 for Windows (Software Statsoft Release 7).

Results

Phytochemical analysis

The rosemary EO was characterized by a total of 23 compounds, accounting for 99.8% of the total composition. The chromatographic results, expressed as gas chromatography peak area percentages and calculated without any response factor correction, highlighted a very high percentage of oxygenated monoterpenes

(51.8%) and hydrocarbon monoterpenes (44.5%), and a lower percentage of hydrocarbon sesquiterpenes (2.7%) and other compounds (0.8%). The main compounds were α -pinene (21.5%), bornyl acetate (16.8%), borneol (10.2%), camphor (9.7%), camphene (7.5%), 1,8-cineole (7.4%), verbenone (4.4%), limonene (4.3%), and β -pinene (3.0%).

Effect of scarification on seed germination

Highly significant differences ($p < 0.001$) were detected among different scarification pretreatments (Fig. 1). In particular, the sandpaper No. 80, the scalpel, and nail clippers were shown to be the best methods, with 100% of germinated seeds in the shortest time ($\text{TI} > 8$, for all). In contrast, the germination percentages obtained by chemical scarification were very low ($<50\%$) independently of the tested immersion time (TI of ca. 0.9).

As the fastest germination times were obtained by scarification using sandpaper No. 80, we used this scarification technique for the following germination tests using EO.

Effects of temperature and EO on seed germination

Preliminary experiments were carried out to evaluate the effect of the individual components in the culture medium. The addition of empty liposomes had no effect on germination, as the final percentages were statistically comparable ($p > 0.05$) with control (ca. $>95\%$ for both). Both the EO concentration (Co) and temperature (Te) had highly significant effect ($p < 0.001$) on the final germination, while population (Po) did not affect the results ($p > 0.05$; Table 2).

Final germination of all the tested populations decreased with increasing EO amount (Fig. 2). In particular, Asal1 showed inhibition of its seed germination at 10 and 20 $^{\circ}\text{C}$ starting from 7.81 μL , while at 15 $^{\circ}\text{C}$ the reduction in final germination occurred only starting from 15.62 μL . The highest tested EO 15.62 $\mu\text{L}/\text{Petri dish}$ concentration caused an inhibition of germination of ca. 85% at 20 $^{\circ}\text{C}$ and of ca. 95% at 10 and 15 $^{\circ}\text{C}$. For Asal2, no significant inhibition ($p > 0.05$) occurred up to 7.81 μL at all temperatures, while total inhibition was observed at 10 $^{\circ}\text{C}$ under 15.62 μL . Asal3 seeds showed a decrease starting from 7.81 μL at both the two tested temperatures (15 and 20 $^{\circ}\text{C}$) and the 15.62 μL amount caused the inhibition of ca. 85% at 15 $^{\circ}\text{C}$ and ca. 90% at 20 $^{\circ}\text{C}$. For Asal4, a significant decrease in seed germination was observed starting from 3.90 μL at 20 $^{\circ}\text{C}$, while starting from 7.81 μL at 15 $^{\circ}\text{C}$. The highest tested EO amount inhibited the final germination of ca. 73% at both the two tested temperatures. Overall, the suboptimal temperatures for this species (10 and 20 $^{\circ}\text{C}$) caused a highly significant effect ($p < 0.001$) of the EO on the seed germination and this pattern was also confirmed by the statistical analyses (interaction $\text{Co} \times \text{Te}$ in Table 2).

Fig. 1. Final germination of *Acacia saligna* seeds under different scarification pretreatments. A Kruskal–Wallis test was carried out to identify differences among the tested pretreatments. Values with different letters are significantly different at $p < 0.05$ (by Mann–Whitney U test).




Table 2. Effect of essential oil concentration (Co), temperature (Te), population (Po), and their interactions on the final germination of *Acacia saligna* seeds from the four investigated populations.

Effect	SS	df	MS	F	p
Co	181245.6	4	45311.4	702.14	***
Te	2250.0	1	2250.0	34.87	***
Po	448.8	3	149.6	2.32	ns
Co × Te	1760.0	4	440.0	6.82	***
Co × Po	1351.2	12	112.6	1.74	ns
Te × Po	214.8	3	71.6	1.11	ns
Co × Te × Po	1195.2	12	99.6	1.54	ns
Error	7744.0	120	64.5		

Note: p values were considered not significantly ($p > 0.05$, ns), or highly significantly ($p < 0.001$, ***) different, by three-way ANOVA. SS, Sum of squares; df, degrees of freedom; MS, mean square; F, Fisher variable.


Discussion and conclusions

This study highlights a qualitative and quantitative chemical composition of the rosemary EO that is the same as other oils used for the management of IAS and weeds, characterized by a phytotoxic effect on the seed germination of several species (Angelini et al. 2003; Atak et al. 2016). It is well known that the EOs are characterized by high concentrations of two or three compounds, compared with the other components (which are present in trace) and generally these major compounds determine the biological proprieties of the EOs (Bakkali et al. 2008). Some monoterpenes present in the rosemary EO, such as α -pinene, limonene,

1,8-cineole, borneol, camphor, are known to inhibit seed germination in some species (De Martino et al. 2012). Based on its chemical composition, we hypothesised that EO would inhibit germination of *Acacia saligna*.

The use of empty liposomes in the culture medium did not influence the final germination of *Acacia saligna*. The incorporation of EO in liposomes limited the normal loss of EO volatile components, thus delivering them into the seed and potentiating EO efficacy. This innovative method, used for the first time, might be used in further studies on species with slow germination because it preserves EO composition. Further studies might investi-

Fig. 2. Effect of different concentrations of *Rosmarinus officinalis* essential oil (EO) on the seed germination of four *Acacia saligna* populations at constant temperatures (10, 15, 20 °C). Bars with different letters are significantly different at $p < 0.05$ (three-way ANOVA followed by Fisher's least significant difference post hoc test). Data are the means ($\pm$ SD) of four replicates. See Table 1 for the explanation of population codes. Tr1, Control (only 1% agar); Tr2, soy lecithin + agar 1%; Tr3, 3.90 μ L EO; Tr4, 7.81 μ L EO; Tr5, 15.62 μ L EO.



gate the use of EO liposomes applied to the shoot of *Acacia saligna* plants, in the field.

Our study provides preliminary information about the phytotoxic effect of rosemary EO on *Acacia saligna* seed germination. In particular, it showed that *Acacia saligna* seeds germinated when exposed to up to 15.62 μ L of *R. officinalis* EO in the substrate, independently from the seed provenance. The interaction between temperature and EO presence in the substrate amplified the inhibitory effect of increasing EO amount. Overall, the inhibi-

tion of the seed germination was low, up to 7.81 μ L, but the highest tested concentration showed that *Acacia saligna* seeds were highly sensitive to this EO, resulting in a potentially effective method for inhibiting germination in *Acacia saligna* that warrants further investigation.

The results obtained in the present study are similar to those of previous studies, which showed the inhibition of the seed germination of several weeds (and cultivated plants associated to them) above a certain concentration of rosemary EO. In particular, as showed by Atak et al.

(2016) for *Avena sterilis* L. seeds, concentrations of rosemary EO from 2 to 16 μL /Petri dish caused a decrease in germination from 61% to 12%, whilst at least 2 μL /Petri dish inhibited germination of *Sinapis arvensis* L. by 89%. In our study, the germination inhibition of *Acacia saligna* seeds was more similar to that observed for *Avena sterilis*, showing that the rosemary EO effects are species-specific.

The present study showed that no *Acacia saligna* seed recovered the ability to germinate after EO exposure followed by either of the two washing methods. This indicates that the rosemary EO might cause irreversible damage to *Acacia saligna* seeds. Witzke et al. (2010) asserted that the phytotoxic effects of EOs on seed germination and seedling development and survival occur through several actions on different functional processes, such as: loss of membrane integrity, ion and solute transport through membranes, synthesis of molecules and macromolecules (e.g., membrane lipids, chlorophyll, proteins, and nucleic acids), and cell division. Moreover, EOs may trigger changes in the mitochondrial adenosine triphosphate production, which compromise the mitochondrial metabolism, in particular during seed germination and early seedling development (Botha et al. 1992; Neuburger et al. 1996). It was not possible to determine which physiological mechanism(s) caused the inhibition of germination of *Acacia saligna*.

The potential for using phytotoxicity in the management of weeds and invasive species has been well documented (Putnam 1988; An et al. 1998). Moreover, germination inhibition by EOs applied to seeds of various crops, weeds, and IAS has been reported in several studies (Nishimura et al. 1982; Azirak and Karaman 2008; Mutlu and Atici 2009; Gitsopoulus et al. 2013).

Recently, EOs obtained from natural indigenous plants were widely tested in the control of weeds and IAS. The potential phytotoxic effect caused by the EOs of indigenous plant species might be a reliable means for the control and management of IAS and weeds.

On the basis of our encouraging results and the verified ability of the rosemary oil to inhibit the seed germination of our invasive study species, the EO of rosemary might be used to control the invasiveness of *Acacia saligna*. In conclusion, these results may be useful for the control against IAS, thus allowing the conservation of native plant species and the preservation of the Mediterranean habitats.

Further studies are in progress to evaluate the potential phytotoxic effect of *R. officinalis* EO on seedling development and survival of *Acacia saligna*, so that these treatments may be applied to reduce its establishment and seedling recruitment directly in natural habitats.

Acknowledgements

The authors thank the Co.S.Me.Se and BG-SAR for financial support and the entire research group, in particular Lina Podda, Maria Letizia Manca, and Carla Caddeo.

Particular thanks also to Anna Rita Succa for the linguistic revision of the manuscript.

References

- Amiot, J., Salmon, Y., Collin, C., and Thompson, J.D. 2005. Differential resistance to freezing and spatial distribution in a chemically polymorphic plant *Thymus vulgaris*. *Ecol. Lett.* 8(4): 370–377. doi:10.1111/j.1461-0248.2005.00728.x.
- An, M., Pratley, J.E., and Haig, T. 1998. Allelopathy: from concept to reality. In *Proceedings of the 9th Australian Agronomy Conference*, Australian Agronomy Society, Wagga Wagga, Australia. pp. 563–566.
- Angelini, L.G., Carpanese, G., Cioni, P.L., Morelli, I., Macchia, M., and Flamini, G. 2003. Essential oils from Mediterranean Lamiaceae as weed germination inhibitors. *J. Agr. Food Chem.* 51(21): 6158–6164. doi:10.1021/jf0210728.
- Atak, M., Mavi, K., and Uremis, I. 2016. Bio-herbicidal effects of oregano and rosemary essential oils on germination and seedling growth of bread wheat cultivars and weeds. *Rom. Biotechnol. Lett.* 21(1): 11149–11159.
- Azad, S., Manik, M.R., Hasan, S., and Matin, A. 2011. Effect of different pre-sowing treatments on seed germination percentage and growth performance of *Acacia auriculiformis*. *J. For. Res.* 22(2): 183–188. doi:10.1007/s11676-011-0147-y.
- Azirak, S., and Karaman, S. 2008. Allelopathic effect of some essential oils and components on germination of weed species. *Acta Agr. Scand. B Soil Plant*, 58(1): 88–92. doi:10.1080/09064710701228353.
- Bakkali, F., Averbeck, S., Averbeck, D., and Idaomar, M. 2008. Biological effects of essential oils - a review. *Food Chem. Toxicol.* 46(2): 446–475. doi:10.1016/j.fct.2007.09.106. PMID: 17996351.
- Bartolucci, F., Peruzzi, L., Galasso, G., Albano, A., Alessandrini, A., Ardenghi, N.M.G., Astuti, G., Bacchetta, G., Ballelli, S., Banfi, E., Barberis, G., Bernardo, L., Bouvet, D., Bovio, M., Cecchi, L., Di Pietro, R., Domina, G., Fascetti, S., Fenu, G., Festi, F., Foggi, B., Gallo, L., Gottschlich, G., Gubellini, L., Iamónico, D., Iberite, M., Jiménez-Mejías, P., Lattanzi, E., Marchetti, D., Martinetto, E., Masin, R.R., Medagli, P., Passalacqua, N.G., Peccenini, S., Pennesi, R., Pierini, B., Poldini, L., Prosser, F., Raimondo, F.M., Roma-Marzio, F., Rosati, L., Santangelo, A., Scoppola, A., Scortegagna, S., Selvaggi, A., Selvi, F., Soldano, A., Stinca, A., Wagensommer, R.P., Wilhelm, T., and Conti, F. 2018. An updated checklist of the vascular flora native to Italy. *Plant Biosyst.* 152(2): 179–303. doi:10.1080/11263504.2017.1419996.
- Bellumori, M., Michelozzi, M., Innocenti, M., Congiu, F., Cencetti, G., and Mulinacci, N. 2015. An innovative approach to the recovery of phenolic compounds and volatile terpenes from the same fresh foliar sample of *Rosmarinus officinalis* L. *Talanta*, 131: 81–87. doi:10.1016/j.talanta.2014.07.073. PMID: 25281076.
- Botha, F.C., Potgieter, G.P., and Botha, A.M. 1992. Respiratory metabolism and gene expression during seed germination. *Plant Growth Regul.* 11(3): 211–224. doi:10.1007/BF00024560.
- Callaway, R.M., and Aschehoug, E.T. 2000. Invasive plants versus their new and old neighbors: a mechanism for exotic invasion. *Science*, 290(5491): 521–523. doi:10.1126/science.290.5491.521. PMID:11039934.
- Chizzola, R. 2006. Volatile compounds from some wild growing aromatic herbs of the Lamiaceae from southern France. *Plant Biosyst.* 140(2): 206–210. doi:10.1080/11263500600756587.
- Correia, M., Castro, S., Ferrero, V., Crisóstomo, J.A., and Rodríguez-Echeverría, S. 2014. Reproductive biology and success of invasive Australian acacias in Portugal. *Bot. J. Linn. Soc.* 174(4): 574–588. doi:10.1111/boj.12155.
- Crisóstomo, J.A., Rodríguez-Echeverría, S., and Freitas, H. 2013.

- Co-introduction of exotic rhizobia to the rhizosphere of the invasive legume *Acacia saligna*, an intercontinental study. *Appl. Soil Ecol.* **64**: 118–126. doi:10.1016/j.apsoil.2012.10.005.
- DAISIE. 2009. Handbook of alien species in Europe. Springer, Dordrecht. [ISBN 978-1-4020-8279-5.] Available by 17 November 2008 from <http://www.europe-aliens.org>; www.springer.com/life+sci/ecology/book/978-1-4020-8279-5 [accessed 30 January 2016].
- Dajić-Stevanović, Z., Šoštarić, I., Marin, P.D., Stojanović, D., and Ristić, M. 2008. Population variability in *Thymus glabrescens* Willd. from Serbia: morphology, anatomy and essential oil composition. *Arch. Biol. Sci.* **60**(3): 475–483. doi:10.2298/ABS0803475D.
- De Martino, L., Mancini, E., Marandino, A., de Almeida, L.F.R., and De Feo, V. 2012. Chemistry and antigerminative activity of essential oils and monoterpenoids from Mediterranean plants. *Curr. Bioact. Compd.* **8**(1): 13–49. doi:10.2174/157340712799828179.
- Del Vecchio, S., Acosta, A., and Stanisci, A. 2013. The impact of *Acacia saligna* invasion on Italian coastal dune EC habitats. *C. R. Biol.* **336**(7): 364–369. doi:10.1016/j.crv.2013.06.004. PMID: 23932256.
- Doran, J., and Turnbull, J. 1997. Australian trees and shrubs: species for land rehabilitation and farm planting in the tropics. 2nd ed. Australian Centre for International Agricultural Research, Canberra, Australia.
- Drew, B.T., Gonzalez-Gallegos, J.G., Xiang, C.-L., Kriebel, R., Drummond, C.P., Walker, J.B., and Sytsma, K.J. 2017. *Salvia* united: the greatest good for the greatest number. *Taxon*, **66**(1): 133–145. doi:10.12705/661.7.
- Fox, J.E.D. 1995. A review of the ecological characteristics of *Acacia saligna* (Labill.) H. Wendl. *Mulga Research Centre Journal*, **12**: 39–56.
- Gitsopoulus, T., Chatzopoulou, P., and Georgoulas, I. 2013. Effects of essential oils of *Lavandula x hybrida* Rev., *Foeniculum vulgare* Mill. and *Thymus capitatus* L. on the germination and radical length of *Triticum aestivum* L., *Hordeum vulgare* L., *Lolium rigidum* L. and *Phalaris brachystachys* L. *J. Essent. Oil-Bear. Plant*, **16**(6): 817–825. doi:10.1080/0972060X.2013.841364.
- Holmes, P.M., and Cowling, R.M. 1997a. The effects of invasion by *Acacia saligna* on the guild structure and regeneration capabilities of South African fynbos shrublands. *J. Appl. Ecol.* **34**(2): 317–332. doi:10.2307/2404879.
- Holmes, P.M., and Cowling, R.M. 1997b. Diversity, composition and guild structure relationships between soil-stored seed banks and mature vegetation in alien plant-invaded South African fynbos shrublands. *Plant Ecol.* **133**(1): 107–122. doi:10.1023/A:1009734026612.
- Holmes, P.M., MacDonald, I.A.W., and Juritz, J. 1987. Effects of clearing treatment on seed banks of the alien invasive shrubs *Acacia saligna* and *Acacia cyclops* in the Southern and South-Western Cape, South Africa. *J. Appl. Ecol.* **24**(3): 1045–1051. doi:10.2307/2404000.
- Iason, G.R., Dicke, M., and Hartley, E. (Editors). 2012. The ecology of plant secondary metabolites: from genes to global processes. Ecological reviews. Cambridge University Press, Cambridge, New York.
- Khan, M.A., and Ungar, I.A. 1984. The effect of salinity and temperature on the germination of polymorphic seeds and growth of *Atriplex triangularis* Willd. *Am. J. Bot.* **71**(4): 481–489. doi:10.1002/j.1537-2197.1984.tb12533.x.
- Kutiel, P.B., Cohen, O., and Shoshany, M. 2004. Invasion rate of the alien species *Acacia saligna* within coastal sand dune habitats in Israel. *Isr. J. Plant Sci.* **52**(2): 115–124. doi:10.1560/8BK5-GFVT-NQ9J-TLN8.
- Le Maitre, D.C., Gaertner, M., Marchante, E., Ens, E.J., Holmes, P.M., Pauchard, A., O'Farrell, P.J., Rogers, A.M., Blanchard, R., Blignaut, J., and Richardson, D.M. 2011. Impacts of invasive Australian acacias: implications for management and restoration. *Divers. Distrib.* **17**: 1015–1029. doi:10.1111/j.1472-4642.2011.00816.x.
- Lorenzo, P., Palomera-Perez, A., Reigosa, M.J., and Gonzalez, L. 2011. Allelopathic interference of invasive *Acacia dealbata* Link on the physiological parameters of native understory species. *Plant Ecol.* **212**(3): 403–412. doi:10.1007/s11258-010-9831-9.
- Meloni, F., Dettori, C.A., Mascia, F., Podda, L., and Bacchetta, G. 2015. What does the germination ecophysiology of the invasive *Acacia saligna* (Labill.) Wendl. (Fabaceae) teach us for its management? *Plant Biosyst.* **149**(2): 242–250. doi:10.1080/11263504.2013.797032.
- Midgley, S.J., and Turnbull, J.W. 2003. Domestication and use of Australian acacias: case studies of five important species. *Aust. Syst. Bot.* **16**(1): 89–102. doi:10.1071/SB01038.
- Mitić, V., Jovanović, O., Stankov-Jovanović, V., Zlatkovic, B., and Stojanovic, G. 2011. Analysis of the Essential Oil of *Teucrium polium* ssp. *capitatum* from the Balkan Peninsula. *Nat. Prod. Commun.* **7**(1): 83–86. PMID:22428254.
- Morris, T.L., Esler, K.J., Barger, N.N., Jacobs, S.M., and Cramer, M.D. 2011. Ecophysiological traits associated with the competitive ability of invasive Australian acacias. *Divers. Distrib.* **17**(5): 898–910. doi:10.1111/j.1472-4642.2011.00802.x.
- Musil, C.F. 1993. Effect of invasive Australian acacias on the regeneration, growth and nutrient chemistry of South African lowland fynbos. *J. Appl. Ecol.* **30**(2): 361–372. doi:10.2307/2404637.
- Mutlu, S., and Atici, Ö. 2009. Allelopathic effect of *Nepeta meyeri* Benth. extract on seed germination and seedling growth of some crop plants. *Acta Physiol. Plant*, **31**(1): 89–93. doi:10.1007/s11738-008-0204-0.
- Neuburger, M., Rébeillé, F., Jourdain, A., Nakamura, S., and Douce, R. 1996. Mitochondria are a major site for folate and thymidylate synthesis in plants. *J. Biol. Chem.* **271**(16): 9466–9472. doi:10.1074/jbc.271.16.9466. PMID:8621617.
- Nishimura, H., Kaku, K., and Mizutani, J. 1982. Allelopathic substances, (±)-p-Menthane-3,8-diols isolated from *Eucalyptus citriodora* Hook. *Agric. Biol. Chem.* **46**(1): 319–320. doi:10.1080/00021369.1982.10865062.
- Orchard, A.E., and Wilson, A.J.G. 2001. Flora of Australia. 11A, Mimosaceae, Acacia part 1. Australian Biological Resources Study, Canberra, Australia.
- Pintore, G., Usai, M., Bradesi, P., Juliano, C., Boatto, G., Tomi, F., Chessa, M., Cerri, R., and Casanova, J. 2002. Chemical composition and antimicrobial activity of *Rosmarinus officinalis* L. oils from Sardinia and Corsica. *Flavour Fragr. J.* **17**(1): 15–19. doi:10.1002/ffj.1022.
- Pirbalouti, A.G., Ghahfarokhi, B.B., Ghahfarokhi, S.A.M., and Malekpoor, F. 2015. Chemical composition of essential oils from the aerial parts and underground parts of Iranian valerian collected from different natural habitats. *Ind. Crops Prod.* **63**: 147–151. doi:10.1016/j.indcrop.2014.10.017.
- Putnam, A.R. 1988. Allelochemicals from plants as herbicides. *Weed Technol.* **2**(4): 510–518. doi:10.1017/S0890037X00032371.
- Regnault-Roger, C., Vincent, C., and Arnason, J.T. 2012. Essential oils in insect control: low-risk products in a high-stakes world. *Annu. Rev. Entomol.* **57**: 405–424. doi:10.1146/annurev-ento-120710-100554. PMID:21942843.
- Rice, E.L. 1984. Allelopathy. Academic Press, University of Minnesota, Minn., USA.
- Richardson, D.M., Carruthers, J., Hui, C., Impson, F.A.C., Miller, J.T., Robertson, M.P., Rouget, M., Le Roux, J.J., and Wilson, J.R.U. 2011. Human-mediated introductions of Australian acacias - a global experiment in biogeography. *Divers. Distrib.* **17**(5): 771–787. doi:10.1111/j.1472-4642.2011.00824.x.
- Santo, A., Mattana, E., and Bacchetta, G. 2015. Inter- and intra-specific variability in seed dormancy loss and germination

- requirements in the *Lavatera triloba* aggregate (Malvaceae). *Plant Ecol. Evol.* **148**(1): 100–110. doi:[10.5091/plevevo.2015.988](https://doi.org/10.5091/plevevo.2015.988).
- Strydom, M., Esler, K.J., and Wood, A.R. 2012. *Acacia saligna* seed banks: sampling methods dynamics, Western Cape, South Africa. *S. Afr. J. Bot.* **79**: 140–147. doi:[10.1016/j.sajb.2011.10.007](https://doi.org/10.1016/j.sajb.2011.10.007).
- Thompson, J.D. 2005. *Plant evolution in the Mediterranean*. Oxford University Press, Oxford, U.K.
- Werner, C., Zumkier, U., Beyschlag, W., and Maguas, C. 2010. High competitiveness of a resource demanding invasive *Acacia* under low resource supply. *Plant Ecol.* **206**(1): 83–96. doi:[10.1007/s11258-009-9625-0](https://doi.org/10.1007/s11258-009-9625-0).
- Witzke, S., Duelund, L., Kongsted, J., Petersen, M., Mouritsen, O.G., and Khandelia, H. 2010. Inclusion of terpenoid plant extracts in lipid bilayers investigated by molecular dynamics simulations. *J. Phys. Chem. B*, **114**(48): 15825–15831. doi:[10.1021/jp108675b](https://doi.org/10.1021/jp108675b). PMID:[21070035](https://pubmed.ncbi.nlm.nih.gov/21070035/).
- Zuzarte, M., Gonçalves, M.J., Cavaleiro, C., Cruz, M.T., Benzarti, A., Marongiu, B., Maxia, A., Piras, A., and Salgueiro, L. 2013. Antifungal and anti-inflammatory potential of *Lavandula stoechas* and *Thymus herba-barona* essential oils. *Ind. Crops Prod.* **44**: 97–103. doi:[10.1016/j.indcrop.2012.11.002](https://doi.org/10.1016/j.indcrop.2012.11.002).